

## REPRODUCTIVE CYCLE AND MARICULTURE POTENTIAL OF THE RABBITFISH *SIGANUS* *CANALICULATUS* IN SAUDI ARABIA

Adel A. Tharwat

Department of Animal Production, Faculty of Agriculture, Cairo  
University, Giza, Egypt.

**Keywords:** Rabbitfish, *Siganus canaliculatus*, reproduction,  
fisheries, Arabian Gulf.

### ABSTRACT

The present work was meant to the reproductive cycle of the rabbitfish *Siganus canaliculatus*, one of the economically important marine fish inhabiting the Arabian Gulf of Saudi Arabia coast. The future expectations of development of *Siganus* mariculture were described. This study revealed the sex ratio, gonads maturity stages, size at first sexual maturity, spawning season, ova-diameter and fecundity. The length at first sexual maturity (Lm50) was found to be 18 cm and 19 cm for males and females, respectively. Fishes larger than 22 cm total length of both sexes were found to be fully mature. Although, the gonado-somatic index (GSI) of *Siganus canaliculatus* differed significantly ( $P > 0.05$ ) between males and females, it exhibited the same trend for both, where GSI reached its maximum values (peaks) of 18.0 and 22.5 during May for males and females, respectively. The spawning season was found to be short and occurred during the period from April to June, with a high peak during May, which means that one spawning season was exhibited. The egg-diameter distribution was significantly variable in the ripe ovaries with largest egg diameters ranging from 0.3 to 0.4 mm, which constitute about 88 % of the total eggs and revealing one peak during May. Absolute fecundity was increased with increasing length, weight and age of the fish. Fecundity exhibited a curvilinear trend or exponential equation with fish length, fish weight and fish age. The mean sex ratio (Males: Females) for the whole year was 1:1.13. It was recommended that capturing of fish smaller than 20 cm total length should be prohibited to protect the spawning stock biomass of rabbitfish *S. canaliculatus* on the long run. Also, many aspects for mariculture potential were included.

## INTRODUCTION

The White Spotted Rabbitfish *Siganus canaliculatus* (Park, 1797), has a local name 'Safi' in the western coast of Arabian Gulf, as it has been popular with excellent marketing and a good price. Moreover, it is relatively easy to rear and thus considered suitable for aquaculture (Lam, 1974; Bryan & Madraisau, 1977; Juario *et al.*, 1985 ; Hara *et al.*, 1986 a & b). There have been attempts to breed and rear two *Siganus* species in the Arabian Peninsula; the marbled spinefoot *Siganus rivulatus* Forsk (Al Thobaity *et al.*, 1984) and the pearl (white) spotted *Siganus canaliculatus* Park (Al-Ghais, 1993). Few studies are available on some biological aspects of *S. canaliculatus* in the Arabian Gulf, which did not cover the life history. So, the present work aimed to investigate the reproductive cycle of this fish including gonad developmental stages, adult sex ratio, gonado-somatic index, timing of spawning season, size and age at first sexual maturity, egg diameters and fecundity. The mariculture potential of *S. canaliculatus* in the Arabian Peninsula was discussed with the available biological information in the literature, which will help to develop fish production in Arabian Gulf on a scientific basis.

## MATERIALS AND METHODS

Specimens of Rabbitfish *Siganus canaliculatus* were collected monthly during the period from January 2002 to December 2002 by professional fishermen using trammel gillnets at Al-Qatif and Al-Jubil coasts of Saudi Arabia on the Arabian Gulf. Fishes were put immediately in crushed ice and transported to the laboratory, where they were measured and investigated. Date of capture, fish total length (mm) and fish total weight (0.1g) were recorded for each fresh specimen. Fishes were dissected to define their sex, gonads maturity stage and fullness of alimentary canal. Fresh gonads were removed and visually inspected for size, colour, vascularization and presence of milt and oocytes. Maturity stages for Siganids species according to Al-Ghais, (1993) and El-Sayed & Bary, (1994) were adopted for the present study. Gutted fish were weighed to the nearest 0.1g and gonads were weighed to the nearest 0.001g, then the ripe ovaries were fixed immediately in 5 % buffered formaldehyde for fecundity and ova diameter investigation. Fish were identified as either immature (with at stage 1 or 2, being small and undeveloped) or mature (with

enlarged gonads, and in case of females, recruiting vitellogenic oocytes were present). For both males and females, frequency data of status of maturity were assembled on length and body weight. The length at which 50 % of fishes reach their sexual maturity was considered as the length at first sexual maturity " $L_{m50}$ " (Pitt, 1970) and was derived from the length frequency distribution of mature males and females. The gonado-somatic index (G.S.I.) was calculated for each specimen using the following equation:

$GSI=100$  [gonads weight (g)/gutted weight (g)], the spawning season was determined by curvilinear of the average monthly GSI for both males and females. Ripening pair of ovaries were fixed in 5 % formaldehyde in glass tubes, with labels indicating date of capture, fish length, fish weight and gonad weight in order to estimate fecundity. From each three subsamples, each approximately 3 % of gonad weight, were removed and weighed to the nearest 0.001 g and the eggs were counted in a Petri dish under a dissecting microscope at a power of 20 X. The absolute fecundity was calculated as the average number of eggs contained in the ovary per individual fish. The numbers of eggs in the three subsamples were averaged (n), and then the total number of eggs in the two ovaries (N) was estimated as follow:  $N = n \times$  [weight of ovaries / weight of sample]. Fish age was based on an age-length key according to Tharwat (2004). Egg diameters were measured monthly using; a research microscope aided with an ocular micrometer at a power of 100 X. Simple regression, standard deviation of mean values for the relationships between estimated fecundity and fish length, fish weight and fish age were computed and the differences between males and females were tested by analysis of variance, according to Schaffer and Elson, (1975) using the program of SAS, (1990).

## RESULTS AND DISCUSSION

### Maturity stages

A macroscopic description of the gonads maturity stages for males and females of *Siganus canaliculatus* are given in Table (1). Macroscopic criteria such as gonad size, shape, texture and color were used to classify the gonad maturity stages in the fish. The main criteria were adopted by many authors (Munro *et al.*, 1973, Hasse *et al.*, 1977, Chan and Chua, 1980, Hara *et al.*, 1986a ; AL-Ghais, 1993). It was found that *S. canaliculatus* have unequal gonad lobes.

where the left lobe of ovary or testis becomes bigger in size throughout maturation cycle than the right lobe when developed. The left lobe is relatively flat and covers the entire left surface of the cavity. The right one is more rounded and elongated, pushing the gut upward. At the same maturity stage, ovaries and testes were relatively similar in appearance and size for *S. canaliculatus*. Undeveloped or juvenile gonads (stage I) were small, cylindrical and occupied a minute portion of the lower body cavity, located over the posterior part of the pelvic fins, and extending posteriority. In April and May all maturity stages were found, thus the spawning season of *S. canaliculatus* extends from April to June with a peak in May. In general, the gonads development of *Siganus* species is affected by several environmental conditions such as: 1) temperature, where the rapid gonadal development usually occurs when the water temperature ranged from 25-30 °C; 2) lunar cycle, where gonads of *S. canaliculatus* mature during new moon; 3) photoperiod (e.g. 18 h light : 6 h dark retards gonadal maturation of *S. canaliculatus*) (Lam and Soh, 1975); and 4) quantity and quality of diet (e.g. females of *S. guttatus* fed with commercial diet containing 43% protein spawned monthly for 11 months; when lecithin, cod liver oil, or both were added to the diet, spawning occurred for at least 4 consecutive months) (Hara *et al.*, 1986a).

### Sex ratio

The length frequency distribution, percentage of males and females of *S. canaliculatus* and the sex ratio with length intervals are shown in Table (2). It was obvious that sex-ratio deviate significantly from 1:1 among the size classes. However, The overall sex ratio (Males: Females) of this species during the year was 1: 1.13 and did not differ significantly from 1:1. The present result agrees with others on the same species in Qatar that was found to be 1:1.09 for males: females (El-Sayed and Bary, 1994) and slightly differ from that of 1.1:1.0 (Wassef and Abdul Hady, 1997) on the same species in Saudi Arabia. On the other hand, there was a predominance of males amongst the smallest size ranging from 14 to 22 cm total length, while females were predominated for the size larger than 22 cm (Table 2). Similar results indicated that female *S. rivulatus* significantly dominated catches along the Syrian coast (Saad & Sabour, 2001). Hashem (1983) indicated a slight dominance of female *S. rivulatus* (1:13) in Red Sea populations but it is unclear whether this ratio was significantly different from 1:1.

### Gonadosomatic Index

Among the widely accepted measures of sexual activity in fishes is the Gonadosomatic index (GSI), which is an indicator commonly used for depicting annual reproductive cycles in fishes (Crim & Glebe, 1990). The monthly variations of the mean GSI of *Siganus canaliculatus* for males and females were graphically represented in Figure (1). It is obvious that G.S.I. values were relatively higher for females than males. The GSI of *Siganus canaliculatus* nearly exhibited the same trend for two sexes, where its average values increased during the period from March to June then sharply decreased during July and returns to dormancy from August to February. The GSI reached its maximum values (peaks) of 18.0 and 22.5 during May for both males and females, respectively. This result indicated that the rabbitfish *S. canaliculatus* in the western coast of Arabian Gulf of Saudi Arabia spawns once a year, and the spawning season extends from April to June. The reproductive cycle and gonad maturation in fishes also differs during the different months of the year. According to this scale, ripe and spawning fishes were abundant in the present study during April through June. Thus, the present results confirmed that the spawning season of *S. canaliculatus* extends through these months of the year and the peak of sexual activity occurs in May. Similar results were obtained on the same species in Qatar (El-Sayed and Bary, 1994) and in Saudi Arabia during April and May when average temperature varied between 22.5 and 25°C (Wassef and Abdul Hady, 1997) However, Al-Ghais, (1993) indicated that GSI of *S. canaliculatus* in the Southern coast of Arabian Gulf in U.A.E. begins to rise in February to a maximum in April, when full ripeness was attained for most fishes. The relatively early spawning in U.A.E. may be attributed to the higher water temperature. A similar trend was also noticed for *S. rivulatus* and *S. luridus* in the Mediterranean Sea, where their spawning occurred in June, when surface water temperature ranged between 24 and 29 °C (Bariche *et al.*, 2003). Although differences in spawning intensity and duration may vary between years, defining a spawning season for a species should be based on a large number of specimens sampled regularly over a relatively long period and be coupled with physical and chemical factors (e.g. water temperature, salinity and luminosity). Previous studies on the reproductive biology of siganid species have shown that water temperature plays a role in the determination of the timing and duration of spawning season (Lam & Soh, 1975; Popper *et*

*al.*,1976; Amin,1985a. b). In Jeddah, Amin (1985a, b) reported that *S. rivulatus* spawned from March to September. However, in the Mediterranean Sea, a spawning season is restricted to June for *S. rivulatus* as reported by Popper & Gundermann (1975) and Bariche *et al.* (2003)

#### **Fish size at onset of sexual maturity**

The progression of the percentage of mature males and females of the rabbitfish *S. canaliculatus* for the total lengths ranging from 15-34 and 15-36 cm respectively is represented in Figure (2). The estimated mean length at first sexual maturity (Lm50) of *S. canaliculatus* was found to be 18 and 19 cm for males and females, respectively. Specimens larger than 22 cm in total length were found to be fully mature for both sexes. The calculated ages at the onset of sexual maturity (tm50) by applying age determination formula of Tharwat (2004) were 1.27 and 1.36 years for males and females, respectively. These results support those of Hasse *et al.* (1977) on *S. canaliculatus* in Palau. They concluded that the fish reaches sexual maturity within the first year of their life span. Spawning occurred from April to June where high temperature in summer appeared to be a limiting factor for the gonad development, reducing the breeding season. Maturity stages of gonads can be successfully applied in the field for getting an approximate idea about the spawning season and its duration. Investigation of the maturity stages of the ovaries and testes for each specimen of *S. canaliculatus* during all months of the year revealed that males attained their first sexual maturity slightly earlier than females. In comparing the size at first sexual maturity (Lm50) of the present *S. canaliculatus* with previous studies, there seems a good agreement. However, El-Sayed and Bary (1994) found that Lm50 was about 17.7 cm for males and 17.2 cm for females at Qatar coasts. Wassef and Abdul Hady, (1997) reported that Lm50 corresponds to 17 and 18 cm in Arabian Gulf at Saudi Arabia coast for males and females, respectively. From the present study it can be recommended that capturing fish smaller than 20 cm total length should be prohibited in order to protect the spawning stock biomass on the long run.

#### **Egg diameter**

When *Siganus* fish reaches maturity, the eggs appear small, spherical, demersal and strongly adhesive (Subandiyono *et al.*, 2000), except in case of *S. argenteus* that are free-floating and non-adhesive

(Lam. 1974). This sticky layer enables the eggs to attach to any type of substrate, whether floating or static, as spawning sites. The egg-diameter frequency distribution in the ripe ovaries of prespawning females of the rabbitfish *S. canaliculatus* and the monthly variation of the egg-diameter are graphically represented in Figures (3 and 4), respectively. The variation of egg diameter distribution within ripe ovaries, suggests a rather short spawning season, in which eggs were released in batches. The ripe eggs ranging from 0.30-0.40 mm and the smaller eggs ranging from 0.10-0.25 both constitute about 88 % and 12 % from the total eggs in the ripe ovaries, respectively (Fig.3). On the other hand, the mean egg-diameters were 0.12 and 0.17 mm in January and February, respectively, then an abrupt increase was observed during the period from March - May to reach its maximum value of 0.40 mm in May. A gradual decrease took place afterwards during the successive months from June to August, with a mean of 0.21 mm, followed by a sharp decrease to reach its minimum value (0.10 mm) in December (Fig. 4). It is obvious that the monthly egg-diameter distribution of *S. canaliculatus* followed nearly a similar pattern to GSI, while the oocytes attain their maximum developmental size just before or during the breeding season (Shenouda, 1988). Therefore, the present result confirms that *S. canaliculatus* has one definite spawning season, which occurs during the period from April to June throughout the year. Estimation of the mean egg diameter in different months gives an idea about the maximum size of eggs and duration of the spawning season, whether short or long. It also shows one peak during May, confirming the previously deduced results, from the GSI values.

### **Fecundity**

Two terms are usually applied in fecundity studies; namely the absolute and relative fecundity where the former denotes the total number of ripe eggs in the ovary, whereas the relative fecundity denotes the number of these eggs per unit length, weight or age of fish.

### **Relationship between fecundity and fish length**

The relationship of the absolute and relative fecundity with the total length of *Siganus canaliculatus* is shown in Table (3). In general, absolute fecundity was higher in larger fish. Female ranging in total length from 17 to 35 cm produces 58925 to 838652 eggs. The

relationship can be represented by the following exponential equation:  $F = 1.910 L^{3.6540}$ ,  $r = 0.979$ , where  $F$  is the absolute fecundity,  $L$  is the total length of fish (cm),  $r$  is the correlation coefficient of the regression. The relative fecundity increased regularly from 3466 to 23961 eggs / cm with a mean of 5645 eggs per cm length increment. In comparing the present results with those of some authors on *S. canaliculatus* in different Arabian Gulf countries, it was found that the absolute fecundity ranged between 650,000 and 1,000,000 eggs per female in U.A.E. (AL-Ghais, 1993), between 225700 and 1,050,000 eggs in Qatar (El-Sayed and Bary, 1994) and from 42253 to more than one million for fish between 17-41 cm total length in Saudi Arabia (Wassef and Abdul Hady, 1997). The relative variation of fecundity could be attributed to the fish size and localities, which is directly affected by fishing pressure among the successive years.

#### **Relationship between fecundity and fish weight**

Table (4) shows the relation of absolute and relative fecundity with fish weight. The results revealed that the absolute fecundity increased with increasing weight of the fish, which increased from 62420 to 983517 eggs for different weight groups that ranged between 75 to 575 gm. The relation between these two variables could be expressed by the following exponential equation:  $F = 180.12 W^{1.3542}$ ,  $r = 0.9985$ , where  $F$  is the absolute fecundity,  $W$  is fish weight (g),  $r$  is the correlation coefficient of the regression. So a high agreement between the mean observed and calculated fecundity for different weight groups (Table 3). On the other hand, the relative fecundity-weight values slightly increased with weight increment, ranging between 832 – 1710 eggs / g with the mean of 1352 eggs per gram.

#### **Relationship between fecundity and fish age**

The average numbers of eggs for the successive age groups of *Siganus canaliculatus* are shown in Table (5). It was found that the number of eggs greatly varied within any age group. This variation is greater than that between fishes of the same length or weight groups. The relationship between absolute fecundity ( $F$ ) and fish age can be expressed by the following exponential equation:  $F = 55267 G^{1.7}$ ,  $r = 0.9795$ , where  $G$  is the fish age by year,  $r$  is the correlation coefficient of the regression. Generally, the mean number of eggs increased, as the fish got older. Thus, it increased from 55173 eggs in fishes of age



group I to 852376 eggs in fishes of age-group V. This indicates that *Siganus canaliculatus* did not reach the senility age and still possesses an economic importance. However, the relative fecundity (eggs/year) in the successive age groups was increasing with older fish (Table 5). The increase in age is usually accompanied by increase in length and weight, so fecundity increases as well. This increase continues until the senility age, at which it stops increasing. The commercial fish species, which has an economic importance, does not reach this age (Nikolsky, 1963). Generally, the present results indicate that *S. canaliculatus* is a highly fecund species and the relation between fecundity and length or weight of a fish may exhibit a nonlinear trend. These results are comparable with those of earlier workers on Siganids (Lam, 1974; Alcalá and Alcazar, 1979; Woodland, 1979; Tseng and Chan, 1982; Al-Ghais, 1993; El-Sayed and Bary, 1994; Wassef and Abdul Hady, 1997).

#### **Prospective potential of rabbitfish mariculture in Saudi Arabia**

As the biological aspects and feeding habits of rabbitfish vary between species, special attention to one or few commercial species that have potential for mariculture is needed. Considerations include the ease of spawning, with or without hormonal treatment. The chosen species should be fecund, fast growing and suited to intensive culture. Based on the above mentioned findings it can be suggested that *S. canaliculatus* is an important species for mariculture in the Arabian Gulf. This fish has potential prospects to be cultured commercially in some countries i.e. Saudi Arabia, Bahrain, Kuwait, United Arab Emirates and Indonesia as they are suited to the local growing conditions and have a recognized delicacy with a moderate market value (Duray, 1990). Other factors which encourage farming of rabbitfish in Saudi Arabia and Arabian Gulf countries are: 1) It is an excellent food with high market value in many countries. 2) The juveniles and adult fish mostly occupy shallow water (Lam, 1974; Popper *et al.*, 1979). Therefore, the fish can be grown using a simple floating cage in the commercial farming, and don't require a deep cage. 3) They inhabit different types of habitat (e.g. coral reef, sandy and rocky bottom (with or without vegetation), lagoons, river estuaries, and mangrove swamps) (Lam, 1974; Popper and Gundermann, 1975; Woodland and Randall, 1979). 4) Sites suited to this type of fish farm are easy to find in the Arabian Gulf. 5) Rabbitfish are able to take an artificial diet (Bwathondi, 1982; Juario

*et al.*, 1985; Hara *et al.*, 1986a; Subandiyono, 1999). 6) Many types of commercial diet for fish have been produced by a large quantity, which encourages mass production, using intensive culture system. 7) The fish can be cultured in monoculture or polyculture system with milkfish (*Chanos chanos*), mullets (*Mugil* and *Liza* spp.) or seabream fishes (Lichatowich and Popper, 1975; Bagarinao, 1986), without affecting growth. 8) Rabbitfish can be used to control the growth of filamentous algae if they are stocked in shrimp ponds and tropical oyster or clam culture (Chen, 1990). 9) The fish spawns easily, whether naturally or by using hormonal treatment (Ayson and Lam, 1993; Subandiyono *et al.*, 1999 & 2000). 10) The fecundity is relatively high, approximately 0.8 million eggs for 400 g fish and 1.2 million for 520 g fish (Popper and Gundermann, 1976). 11) Though the larvae are very fragile, they can be transported for 2 days using simple equipment (Basyari *et al.*, 1988). 12) Large numbers of rabbitfish juveniles can be collected from coastal waters during certain seasons (Lam, 1974). 13) Labor cost is relatively cheaper. 14) They can be used as a bait to catch tuna (Duray, 1990). On the other hand, there are some undesirable characteristics that included: 1) The larvae have a relatively small mouth gape at the first opening, i.e. about 125  $\mu\text{m}$  (Duray and Kohno, 1988). Therefore, they need small size feeds. 2) The fish are difficult to handle due to the poisonous spines that may induce severe headaches (Herzberg, 1973). 3) The sex is difficult to distinguish except during the breeding season (Duray, 1990). 4) It is difficult to differentiate between the species due to only few morphological differences. Therefore, the identification relies only on the coloration of live fish, habitat, and behavioral characteristics (Woodland and Randall, 1979). In general, rabbitfish culture has a tremendous potential in Saudi Arabia and other Arabian countries.

## REFERENCES

- Alcala, A. C. and Alcazar, S. N. (1979). A study on gonad morphology, oocyte development, gonad index and fecundity in the rabbitfish *Siganus canaliculatus* (Park). *Silliman J.*, 26 (2-3): 147-162.
- Al-Ghais, S. M. (1993). Some aspects of the biology of *Siganus canaliculatus* in the southern Arabian Gulf. *Bull. Mar. Sci.*, 52(3): 886-897.

- Al Thobaity, S. ; Bokhary, F. and Badawi, A. (1984). Cultivation of siganids in the Red Sea. Report Min. Agric. Water Res. Jeddah, Saudi Arabia, 71 pp. (in Arabic).
- Amin, E. M. (1985a). Seasonal developmental changes in the ovaries of *Siganus rivulatus* from the Red Sea. Bull. Inst. Oceanogr. & Fish. (A.R.E.) 11: 131-147.
- Amin, E. M. (1985b). Reproductive cycle of male *Siganus rivulatus* Forsk. with indication to gonadosomatic and hepatosomatic indices. Bull. Inst. Oceanogr. & Fish. (ARE) 11: 149-164.
- Ayson, F. G. and Lam, T. J. (1993). Thyroxine injection of female rabbitfish *Siganus guttatus* broodstock: changes in thyroid hormone levels in plasma, eggs and yolk-sac larvae and its effect on larval growth and survival. Aquacult., 109: 83-93.
- Bagarinao, T. (1986). Yolk resorption, onset of feeding and survival potential of larvae of three tropical marine fish species reared in the hatchery. Mar. Biol., 91: 449-459.
- Bariche, M. ; Harmelin, M. V. and Quignard, J. P. (2003). Reproductive cycle and spawning periods of two Lessepsian siganid fishes on the Lebanese coast. J. Fish Biol., 62: 129-142.
- Basyari, A. ; Danakusumah, E. ; Philip, T. I. ; Pramu, S. and Mustahal dan Isra, M. (1988). Budidaya ikan beronang (*Siganus* sp.). INFIS. No. 60, Direktorat Jenderal Perikanan. 31 hal.
- Bryan, P. G. and Madraisau, B. B. (1977). Larval rearing and development of *Siganus lineatus* (Pisces, Siganidae) from hatching through metamorphosis. Aquacult., 10(3): 243-252.
- Bwathondi, P. O. J. (1982). Preliminary investigations on rabbitfish, *Siganus canaliculatus*, cultivations in Tanzania. Aquacult., 27(2): 205 - 210.

- Chan, E. H. and Chua, T. E. (1980). Reproduction in the greenback grey mullet, *Liza subviridis* (Valenciennes, 1836). *J. Fish Biol.* 16: 505-510.
- Chen, L. C. (1990). *Aquaculture in Taiwan*. Fishing News Books, Oxford, USA. 273pp.
- Crim, L. W. and Glebe, B. D. (1990). Reproduction. In *Methods for Fish Biology* (Schreck, C. B. and Moyle, P. B., eds), pp. 529-553. Bethesda, MD: American Fisheries Society.
- Duray, M. N. (1990). *Biology and culture of siganids*. Aquaculture Department, SEAFDEC, Philippines. 47 pp.
- Duray, M. and Kohno, H. (1988). Effects of continuous lighting on growth and survival of first-feeding larval rabbitfish, *Siganus guttatus*. *Aquacult.*, 72: 73-79.
- El Sayed, A. M. and Bary, K. A. (1994). Life cycle and fecundity of rabbitfish *Siganus canaliculatus* (Teleostei : Siganidae) in the Arabian Gulf. *Oebalia*, 20: 79-88.
- Hara, S. ; Duray, M. ; Parazo, M. and Taki, Y. (1986a). Year-round spawning and seed production of the rabbitfish, *Siganus guttatus*. *Aquacult.*, 59: 259-272.
- Hara, S. ; Kohno, H. and Taki, Y. (1986b). Spawning behavior and early life history of *Siganus guttatus* in the laboratory. *Aquacult.*, 59: 273-285.
- Hashem, M. T. (1983). Biological studies on *Siganus rivulatus* (Forssk.) in the Red Sea. *J. Fac. Mar. Sci.*, 3: 119-127.
- Hasse, J. J. ; Madraisau, B. B. and McVey J. P. (1977). Some aspects of the life history of *Siganus canaliculatus* Park (Pisces: Siganidae) in Palau. *Micronesica*, 13: 297-312.
- Herzberg, A. (1973). Toxicity of *Siganus luridus* (Ruppell) on the Mediterranean coast of Israel. *Aquacult.*, 47: 53-59.

- Juario, J. V. ; Duray, M. N. ; Duray, V. M. ; Nacario, J. F. and Almendras, J. M. E. (1985). Breeding and larval rearing of the rabbitfish *Siganus guttatus* (Bloch). *Aquacult.*, 44: 91-101.
- Lam, T. J. (1974). Siganids: their biology and mariculture potential. *Aquacult.*, 3: 325-354.
- Lam, T. J. and Soh, C. L. (1975). Effect of photoperiod on gonadal maturation in the rabbitfish *Siganus canaliculatus* (Park, 1797). *Aquacult.*, 5: 407-410.
- Lichatowich, T. and Popper, D. (1975). Report on the growth of rabbitfish in fish ponds in Fiji. *Aquacult.*, 5: 211-212.
- Munro, J. L. ; Gaut, V. C. ; Thompson, R. and Reeson, P.H. (1973). The spawning season of Caribbean reef fishes. *J. Fish Biol.*, 5: 69-84.
- Nikolsky, G. V. (1963). The ecology of fishes. Academic Press, London and New York.
- Pitt, T. K. (1970). Distribution, abundance and spawning of yellow tail flounder *Limanda ferruginea* in the new foundland area, the North-West Atlantic. *J. Fish. Res. Board, Canada*, 27 (12): 2261 - 2271.
- Popper, D. and Gundermann, N. (1975). Some ecological and behavioral aspects of siganid populations in the Red Sea and Mediterranean coast of Israel in relation to their suitability for aquaculture. *Aquacult.*, 6: 127-141.
- Popper, D. and Gundermann, N. (1976). A successful spawning and hatching of *Siganus vermiculatus* under field conditions. *Aquacult.*, 7: 291-292.
- Popper, D. ; May, R. C. and Lichatowich, T. (1976). An experiment in rearing larval *Siganus vermiculatus* (Valenciennes) and some observations on its spawning cycle. *Aquacult.*, 7: 281-290.

- Popper, D. H. ; Pitt, R. and Zohar, Y. (1979). Experiments on the propagation of Red Sea siganids and some notes on their reproduction in nature. *Aquacult.*, 16:177-181.
- Saad, A. and Sabour, W. (2001). Donnees preliminaires sur la reproduction de *Siganus rivulatus* (Forsk. 1777). Pisces. Siganidae), dans les eaux cotieres de Syrie. Rapport du 36 Congres de la Commission Internationale Pour l'Exploration Scientifique de la Mer Mediterranee 36 : 319.
- SAS, (1990). Statistical analytical systems user's guide (Vol. 2). Cary, NC : SAS Institute Inc. USA.
- Schaffer, W. M. and Elson, F. P. (1975). The adaptive significance of variations in life history among local populations of Atlantic salmon in North America. *Ecology*, 56: 577-590.
- Shenouda, T. S. (1988). Spawning peculiarities of *Trachurus trachurus* L. (Fam. Carangidae) in the Mediterranean Sea. *Delta J. Sci.*, 12(4): 1711 - 1723.
- Subandiyono, (1999). Growth of rabbitfish, *Siganus* sp., in captivity fed by diets containing different level of soy-lecithin. *Journal of Coastal Development*, 2(3): 419-425.
- Subandiyono, Kokarkin, C. and Dan Hastuti, S. (1999). Paket teknologi formulasi pakan induk ikan beronang (*Siganus* sp.) guna meningkatkan kualitas telur. Tahun II. Lemlit-Universitas Diponegoro, 81 hal.
- Subandiyono, Kokarkin, C. and Dan Hastuti, S. (2000). Paket teknologi formulasi pakan induk ikan beronang (*Siganus* sp.) guna meningkatkan kualitas telur. Tahun III. Lemlit-Universitas Diponegoro. 102 hal.
- Tharwat, A. A. (2004). Population dynamics of the rabbitfish *Siganus canaliculatus* from the Arabian Gulf of Saudi Arabia. Egypt. *J. Aquat. Biol. & Fish.*, (in publication).

REPRODUCTIVE CYCLE  
OF THE *S. CANALICULATUS* IN SAUDI ARABIA

---

137

- Tseng, W. Y. and Chan, K. L. (1982). The reproductive biology of the rabbitfish in Hong Kong. *J. World Maricult. Soc.*, 13: 313-321.
- Wassef, E. A. and Abdul Hady, H. A. (1997). Breeding biology of rabbitfish *Siganus canaliculatus* Park in mid Arabia Gulf. *Fish. Res.*, Amsterdam, 33: 159-166.
- Woodland, D. J. (1979). Rabbitfishes neglected in Australia are important food fish in tropical countries. *Aust. Fish.*, 38(6): 21-23.
- Woodland, D. J. and Randall, J. E. (1979). *Siganus puelloides*, a new species of rabbitfish from the Indian Ocean. *Copeia*, 3: 390-393.

Table (1). The maturity stages of *Siganus canaliculatus* inhabiting the Arabian Gulf.

Maturity stage (Classification)	Description
0 (Immature)	No easily visible gonads in body cavity. Sex cannot be determined macroscopically
I (Maturing)	Gonads are represented by two thin tubular units threadlike, transparent and can be distinguished into testis or ovary.
II (Early developed)	Ovaries begin to swell, enlarged, still translucent and easily recognized. Oocytes cannot visually discernible. Male testes are enlarged, opaque and easily recognized.
III (Mid developed)	Ovaries larger, occupying about 50 % of the body cavity, visible asymmetric and borders are more rounded than testes, with a circular cross-section. Ovaries pale yellow to dark-pink in color. Vascularization starts to appear and Oocytes can visually discernible. Testes marked asymmetry with triangular cross-section. Whitish gray in color and sometimes bluish and Vascularization is not visible.
IV (Ripe)	Enlarged ovaries, solid and occupying about 75% of the body cavity, with rounded borders, marked asymmetric and well-developed vascularization. Colour ranges from pale to darker yellow. Eggs fully formed, numerous and clearly visible to naked eye. Testes usually white, cleared asymmetric with triangular cross-section and some sperms expelled from core when testis is cut.
V (Spawning)	Fish about to spawn. Gonad turgid and filling the body cavity. Swollen ovaries with a dark yellow and sometimes orange colour. Strongly asymmetric and vascularization is well developed. Eggs are transparent and are expelled with a slight pressure on the abdomen. Testes are strongly asymmetric and ivory colour. Vascularization is not clear visible. Characterized by the presence of milt flowing freely when the abdomen is compressed.
VI (Spent)	Ovaries significantly reduced in volume of the body cavity, flaccid and shrink but not fully empty. Ovaries reddish yellow in colour and eggs seen by naked eye. Vascularization relatively well developed. Spent testes relatively hard to recognize. Sometimes little sperm flows. Relatively similar to stage III but more flaccid, dull-gray color, soft, and fleshy in its appearance.



**REPRODUCTIVE CYCLE  
OF THE *S. CANALICULATUS* IN SAUDI ARABIA**

139

**Table (2).** The percentage of males and females of the rabbitfish *Siganus canaliculatus* in length intervals and the sex ratio (Males: Females).

Total length (cm)	Freq.	Males		Females		Sex Ratio (M : F)
		Number	%	Number	%	
14-16	68	38	55.9	30	44.1	1: 0.79
16-18	131	70	53.4	61	46.6	1: 0.87
18-20	234	127	54.3	107	45.7	1: 0.84
20-22	391	197	50.4	194	49.6	1: 0.98
22-24	304	131	43.1	173	56.9	1: 1.32
24-26	209	98	46.9	111	53.1	1: 1.13
26-28	143	64	44.8	79	55.2	1: 1.23
28-30	86	38	44.2	48	55.8	1: 1.26
30-32	36	13	36.00	23	64.0	1: 1.77
32-34	19	3	15.8	16	84.2	1: 5.33
34-36	4	-	-	4	100.0	-
<b>Total</b>	<b>1625</b>	<b>779</b>	<b>47.9</b>	<b>846</b>	<b>52.1</b>	<b>1 : 1.13</b>

**Table (3):** The relationship between fecundity and the total length (L) of the rabbitfish *Siganus canaliculatus* collected from the Arabian Gulf in Saudi Arabia.

Total length (cm)	Mean length (cm)	No. of fish	Absolute fecundity		Relative fecundity (Eggs /-cm)
			observed	calculated	
16-18	17	3	58925	59854	3466
18-20	19	5	88979	89867	4683
20-22	21	4	130251	129546	6202
22-24	23	5	181124	180629	7875
24-26	25	7	250076	244966	10003
26-28	27	6	329815	324516	12215
28-30	29	8	421630	421343	14539
30-32	31	6	537871	537612	17351
32-34	33	4	680253	675590	20614
34-36	35	3	838652	837641	23961
<b>Total</b>	-	<b>51</b>	-	-	-
<b>Mean</b>	<b>26</b>	<b>5.1</b>	<b>214275</b>	<b>214312</b>	<b>5645</b>

Table (4): The relationship between fecundity and the total weight (W) of the rabbitfish *Siganus canaliculatus* collected from the Arabian Gulf in Saudi Arabia.

Total weight (g)	Mean weight (g)	No. of fish	Absolute fecundity		Relative fecundity (Eggs / g)
			observed	calculated	
50-100	75	3	62420	62340	832
100-150	125	5	124279	124508	992
150-200	175	4	196573	196374	1123
200-250	225	5	275967	275986	1227
250-300	275	7	362354	362165	1318
300-350	325	6	455201	454103	1401
350-400	375	7	551703	551207	1471
400-450	425	6	654010	653020	1539
450-500	475	3	760125	759172	1600
500-550	525	3	870682	869364	1658
550-600	575	2	983517	983341	1710
Total	-	51	-	-	-
Mean	325	5.0	531676	531053	1352

Table (5): The relationship between fecundity and the age (G) of the rabbitfish *Siganus canaliculatus* collected from the Arabian Gulf in Saudi Arabia.

Age (year)	No. of fish	Absolute fecundity		Relative fecundity (Eggs / year)	
		observed	calculated	observed	calculated
1	7	55173	55267	55173	55267
2	15	179204	179563	89602	89781
3	17	358725	357744	119575	119248
4	9	581930	583402	145482	145850
5	3	852376	852540	170475	170508
Total	51	-	-	-	-
Mean	10.2	405481	405703	945346	-




Fig. 1 Monthly variations of Gonadosomatic index (GSI, %) for males and females of *Siganus canaliculatus*




Fig. 2 The percentages of mature males and females of *Siganus canaliculatus* showing the length at first spawning (Lm 50).




Fig. 3. Frequency distribution as a percentage (%) of ova diameter in ripe ovaries of the rabbitfish *Siganus canaliculatus* during the spawning season.




Fig. 4. The mean value of egg-diameter distributed monthly for the rabbitfish *S. canaliculatus* collected from the Arabian Gulf.