

Management and Reproduction of the Male Brushtooth Lizardfish *Saurida undosquamis* (Richardson) from Gulf of Suez, Egypt

Amal M. Amin, Magdy M. EL-Halfawy and Amal M. Ramadan
National Institute of Oceanography and Fisheries
P.O.Box: 182, Suez, Egypt

ABSTRACT

Age, growth and reproduction of *Saurida undosquamis* collected from Attaka fishing port from October 2004 to May 2005 was studied by examining 390 specimens caught by trawlers in Suez Gulf waters. The total length of males ranged between 9.5 cm to 29.8 cm. The length (L) – weight (W) relationship was estimated as $W = 0.0042 \cdot L^{3.1315}$. The age data derived from the length frequency data analyzed using the FISAT soft ware were used to calculate the growth parameters of the von Bertalanffy equation (1938). The estimated parameters were $L_{\infty} = 31.03$ cm, $K = 0.44$ and $t_0 = -1.059$. The maximum age was 5 years. The 2-year age group was dominant in the catch. Macroscopic appearance of the testes demonstrated seven stages of maturity as virgin, maturing virgin, developing, developed, gravid, spawning and spent. Males recorded first sexual maturity at a total length 17.4 cm. Gonadosomatic index illustrated that this fish have prolonged spawning season nearly all the year round with main peak in May (2.6). The annual rates of total, natural and fishing mortality were calculated as 1.59, 1.32 and 0.27 year^{-1} respectively. Exploitation rate E was estimated as 0.83. The relative yield per recruit was estimated using the parameters obtained for male. The results suggest that the present level of fishing and natural mortality are much higher than that gives the maximum yield per recruit.

Keywords: Gulf of Suez, male *Saurida undosquamis*, age, growth, first sexual maturity, reproductive biology, exploitation rates, yield per recruit.

INTRODUCTION

The present study, which deals with the management and reproduction of *Saurida undosquamis* (males) Richardson from the Gulf of Suez, Egypt is the second contribution in the study of growth and reproduction of this species.

Brushtooth lizardfish *Saurida undosquamis* is one of the most important commercial and economic fish species of family Synodontidae from Gulf of Suez. Family Synodontidae is represented in the trawl catch by five species; *Saurida tumbil*, *Saurida undosquamis*, *Saurida longimanus*, *Synodus hoshinonis* and *Trachinocephalus myops*. In the Gulf of Suez, *Saurida tumbil* is the most important constitute; about 65% of the Lizard fish, followed by *S. undosquamis* which contributes 33% of the catch, while the other three species are the least

abundant; appear occasionally in the catch (EL-Ganainy, 1997). The dynamics and reproductive biology of *Saurida undosquamis* have been studied in different localities (Rao, 1983; El-Ganainy 1992, 1997 & 2002 and Ramadan 1995 in the Gulf of Suez, Golani, 1993, Abdallah, 2002 and EL-Gresiy, 2005 in Mediterranean. Several studies of this species have been undertaken in localities other than Egypt: such as Siripakhavanich (1990) in Gulf of Thailand; Boonwanich (1991), Erzini (1991), Sousa (1992), Federizon (1993), Ismen (2003) in Iskenderun Bay.

The present study deals with estimating the basic parameters required for assessing the status of *Saurida undosquamis* male from the Gulf of Suez in relation to gonad development and this information would help in the proper management of the Gulf of Suez trawl fishery and in the achievement of its optimum sustainable yield.

MATERIAL AND METHODS

Samples (390 fish ranging from 9.5 – 29.8cm TL) were collected from the Attaka harbour during the period from October 2004 to May 2005. The total length to the nearest millimeter, Samples were measured to study the length frequency distribution which divided into 1.0 cm length class. The relation between the total length (L) and total weight (W) was computed using the formula; $W = a L^b$ where a and b are constants whose values were estimated by the least square method.

The FAO-ICLARM Stock Assessment Tools (FiSAT; Gayanilo *et al.* 1997) was used for all computation, aging determined by Battacharya (1967) method which depended on analysis of length frequencies, the growth parameters (L_{∞} , K and t_0) of *Saurida undosquamis* females. The growth performance index (ϕ) was computed according to the formula of Pauly and Munro (1984) as: $\phi = \text{Log } K + 2\text{Log } L_{\infty}$.

The sex and maturity stage of each specimen were determined by visual and microscopic examination of the gonads. The stages of maturation were classified according to Holden and Raitt's (1974) scale. The gonado somatic index (GSI) was calculated monthly by the equation:

$$\text{GSI} = (\text{gonad weight/fish weight without gonad}) * 100$$

The total mortality coefficient "Z" was estimated using the method of Pauly (1983a). The natural mortality coefficient "M" was estimated using the formula of Pauly (1980), while the fishing mortality coefficient "F" was estimated as: $F = Z - M$.

The exploitation rate "E" was calculated using the formula of Gulland(1971) as: $E = F/Z$.

The length at first capture " L_c " was estimated by the analysis of catch curve using the method of Pauly (1984a&b).

Several analytical models are used for the estimated of yield per recruit. The most often used model is that of Beverton and Holt (1957). The model is based on the calculation of yield per recruit under a particular set of fishing mortality and age at first capture. The model can be written in the form suggested by Gulland (1969) as follows:

$$\frac{Y}{R} = Fe^{-M(t_c-t_r)}W_{\infty}\left(\frac{1}{Z} - \frac{3S}{Z+K} + \frac{3S^2}{Z+2K} - \frac{S^3}{Z+3K}\right)$$

where:-

Y/R = yield per recruit

T_c = age at first capture.

T_r = age at recruitment.

Where

$$S = e^{-K(T_c-t_0)}$$

K = von Bertalanffy growth parameter.

W_{∞} = asymptotic body weight

M = natural mortality coefficient.

The age at first capture (T_c) was calculated by converting the length at first capture (L_c) to age by means of the von Bertalanffy growth equation:

$$T_c = -\frac{1}{K} \ln\left(1 - \left(\frac{L_c}{L_{\infty}}\right)^3\right) + t_0$$

(Tr) was computed by the conversion of the length at recruitment (Lr) using the following version of the von Bertalanffy formula as follows:

$$T_r = -\frac{1}{K \ln \left[1 - \left(\frac{L_r}{L_\infty} \right) \right]} + t_0$$

RESULTS AND DISCUSSION

I- Age Determination

Age of male *Saurida undosquamis* in the Gulf of Suez was estimated according to the analysis of length-frequencies data using the Bhattacharya (1967) method. The obtained results revealed that the maximum life span of *Saurida undosquamis* males is five years at length of 13.99, 20.24, 24.13, 26.01 and 28.05 cm. (Fig. 1). This agrees with the finding of EL-Ganainy (2004).




Figure 1: Bhattacharya method of *Saurida undosquamis* males from the Gulf of Suez.

II- Length - Weight Relationship

Length and weight measurements of 390 specimens were used to describe the length-weight relationship of *Saurida undosquamis* males in the Gulf of Suez (Figure 2). Their total lengths varied between 9.5 and 29.8 cm, while the total weights ranged between 17.4 and 160.5 g. The obtained equation was as follow:

$$W = 0.0042 L^{3.1315} \quad (r = .985)$$




Figure 2: Length-weight relationship of *Saurida undosquamis* males from the Gulf of Suez.

III- Growth Parameters

The von Bertalanffy growth parameters (L_{∞} , K and t_0) were estimated and the obtained equations were as follow:

$$\text{For growth in length } L_t = 31.03 (1 - e^{-0.4(t+1.3)})$$

$$\text{For growth in weight } W_t = 197.14 (1 - e^{-0.4(t+1.3)})^{3.1315}$$

IV- Growth Performance Index (ϕ)

The growth performance index (ϕ) of *Saurida undosquamis* males was computed as 2.63.

V- Mortality Rates

The results from Figure (3) indicate that the total mortality coefficient "Z" was estimated as 1.59 year^{-1} . The value of natural mortality coefficient "M" was estimated as 0.27 year^{-1} , while fishing mortality coefficient "F" was estimated as 1.32 year.




Figure 3: Estimation of Z of *Saurida undosquamis* males from the Gulf of Suez.

VI- Exploitation Rate "E"

Exploitation rate "E" was computed using the formula of Gulland (1971) and the obtained E was 0.83. Gulland suggested that the optimum exploitation rate is about 0.5, so the high value of the present exploitation rate indicated that the stock of *Saurida undosquamis* males is overexploited and to maintain this valuable fish resource, the exploitation rate should be reduced below the optimum value as well as increase the length at first capture to be about 20cm.

VII- Length and age at first capture " L_c "

The length at first capture was obtained as $L_{50\%} = 15.79\text{cm}$, which corresponded to an age of 1.81 (Fig. 4).




Figure 4: Length at first capture L_c of *Saurida undosquamis* males from the Gulf of Suez.

Stage VI (Spawning): Testes are slightly shrunk and flaccid but not completely hollow due to discharge of milt during the spawning process. The testes are white in colour.

Stage VII (Spent): Testes are reduced in size. They are shrunken and collapsed. They are deep white colour.

This result coincides with Latife and Shenoda (1973), Ramadan (1995) and EL-Greisy (2005).

IX- Length at first sexual maturity " L_{50} "

Size at first sexual maturity helps to determine the minimum size that must be avoided in order to protect an adequate spawning stock and ensure at least one spawning for the mature individual. The percentage study divided length of male *S. undosquamis* to different groups and classified them into two main categories; Immature (fish of stages I & II) and mature (stages from III to VII) individuals. Mature individuals of male *S. undosquamis* reach first sexual maturity with percentage 50 % at 17.4 cm. Figure (5) demonstrated that all males with total length less than 15 cm are immature, while other fish longer than 21 cm are mature.




Figure 5: Length at first sexual maturity for male *Saurida undosquamis* collected from gulf of Suez, Red Sea, during the period from October 2004 to June 2005.

Faltas (1993) recorded a range of (15-19 cm) for the same species. Latife and Shenoda (1973) recorded (16 cm) in the Gulf of Suez. This was indicated through studies, which were taken in other regions in the same species such as Budnichenko & Dimitrova, 1979 recorded (12 - 13 cm) in the Arabian sea. These different data from different regions means that length at first sexual maturity are related to the environmental conditions.

X- Gonado somatic index (GSI)

Monthly fluctuations of GSI values are shown in Fig. (6). GSI values range between a minimum value 0.64 in November and a maximum one 2.6 in May. The spawning period showed different peaks but the main one was demonstrated in May. This means that this fish is prolonged spawning. This results agree with Ramadan (1995) and EL-Greisy (2005), who indicated that the main GSI occurs in May although the spawning stages found all the year around.




Figure 6: Monthly variation of the average Gonado-Somatic Index of male *Saurida undosquamis* from gulf of Suez, Red Sea, during the period from October 2004 to May 2005.

XI- Yield per recruit

The yield per recruit of *Saurida undosquamis* collected from the Gulf of Suez during October 2004 to May 2005 was estimated by using Beverton and Holt (1957) model. The following parameters values were used in the calculation:

$$L_{\infty}=31.03 \text{ cm} \quad W_{\infty}=198.49 \quad K=0.395 \text{ year}^{-1} \quad t_0=-0.31 \quad L_r=9.5 \text{ cm}$$

$$L_c=\text{variable} \quad M=0.27 \text{ year}^{-1} \quad Z=1.59 \text{ year}^{-1} \quad F=\text{variable}$$

The results are graphically represented in Fig. (7). From the graph it is evident that, the yield per recruit is zero when the fishing mortality is zero, then the yield per recruit increases rapidly as the fishing mortality increases and reached its maximum value at fishing mortality coefficient of (0.5)after which the yield per recruit decreases with further increasing of fishing mortality. Figure show also that at the present level of fishing mortality ($F= 1.32$ per/year), age at first capture ($T_c= 1.306$ year) and natural mortality coefficient ($M= 0.27$ per/year) the yield per recruit was estimated to be(33.04) gm. This mean that ,the present level of fishing mortality is higher than that which gives the maximum yield per recruit and to obtain the maximum yield per recruit 37.16gm, the fishing mortality coefficient must be reduced from 1.32 to 0.5(62.12%).

To determine the most appropriate age at first capture " T_c " of *Saurida undosquamis* from the Gulf of Suez, which is related to the estimation of the optimum mesh size, the yield per recruit was calculated by applying different values of T_c (2.306and 3.306 years with the present value 1.306 year).the results indicate that with the increasing of T_c a higher yield per recruit can be obtained. It is obvious also that if T_c is 2.306 instead of 1.306 year a maximum yield per recruit of 44.13gm can be obtained at fishing mortality of 1.1 which is close to the present level ($F=1.32$). It is notes also that at $T_c \approx 3.306$ years a maximum

yield per recruit of 47.19gm is obtained. This mean that, the present level of T_c is not the optimum T_c of this fish species in the Gulf of Suez and it must be increase.




Figure 7: Yield per recruit (gm.) of *Saurida undosquamis* (males) from the Gulf of Suez as a function fishing mortality and age at first capture.

To study the variation in the yield per recruit according to the change of the natural mortality coefficient “M”, the yield per recruit of *Saurida undosquamis* was estimated by using different values of “M” and the obtained results represented graphically in Fig. (8) which clear that the yield per recruit decreases with the increase of the natural mortality coefficient.

It is also evident that at the present level of the fishing mortality coefficient ($F=1.32$) and age at first capture ($T_c=1.306$ year) a higher yield per recruit can be obtained (37.89 instead of 33.04gm) when the natural mortality coefficient decreased (0.19 instead of 0.27) if the natural mortality coefficient is higher than the present level (0.35 instead of 0.27) a lower value of yield per recruit was obtained (28.86gm instead of 37.98gm) (Fig. 8).

This means that the yield per recruit decreases with the increasing of natural mortality coefficient and the habitat conservation can increase the yield per recruit through the protection of the nursery ground as well as the protection from pollution and illegal fishing.




Figure 8: Yield per recruit (gm.) of *Saurida undosquamis* (males) from the Gulf of Suez as a function fishing mortality and natural mortality.

The obtained results are in a good agreement with the finding of Sanders *et al.* (1984). They stated that the stock of *Saurida undosquamis* in the Gulf of Suez is fully exploited and any addition in the fishing effort will be associated with a decrease in the catch.

REFERENCES

- Abdallah, M. (2002). Length-weight relationship of fishes caught by trawl off Alexandria, Egypt. *Naga ICLARM Q.*, 25(1):19-20.
- Battacharya, C. G. (1967). A simple method of resolution of a distribution into Gaussian components. *Biometrics*, 23:115-135.
- Beverton, R. J. H. and Holt, S. J. (1957). On the dynamics of exploited fish population. U.K. Min. Agr. Fish. Food, *Ish-Inuest*, 19: 533 pp.
- Boonwanich, T. (1991). Population dynamics of *Saurida elongata* and *S. undosquamis* (Synodontidae) in the southern Gulf of Thailand. *Fishbyte*, 9(1):23-27.
- Budnichenko V. A. and Dimitrova O. S. (1979). Reproductive biology of *Saurida undosquamis* and *Saurida tumbil* (pisces, synodontidae) in the Arabian Sea. *J. Ichthyol.*, 19 (5): 860-867.

- El-Ganainy, A. A. (1992). Biological studies on lizard fishes *Saurida undosquamis* (Pisces, Synodontidae) from the Gulf of Suez. M. Sc. Thesis, Fac. Sci., Ain Shams University, Cairo. Egypt.
- El-Ganainy, A. A. (1997). Population Dynamics of Lizard Fishes (Synodontidae) from the Red Sea. Ph.D. Thesis, Fac. Sci., Suez Canal University, Ismailia. Egypt.
- El-Ganainy, A. A. (2002). "Assessment of the lizard fish *Saurida undosquamis* (Richardson) fishery in the Gulf of Suez, Red Sea." Bull. Nat.Inst. Oceanogr. Fish., ARE., 28 :
- El-Ganainy, A. A. (2004). " Biological characteristic and fishery assessment of the lizard fishes *Saurida undosquamis* from the Red Sea, Egypt ".Egypt. J. Aquatic. Biol. & Fish. , 8 (2): 93-113.
- El-Greisy Z. A. (2005). Reproductive Biology and Histology of Female Brushtooth Lizardfish *Saurida undosquamis* (Richardson), Family: Synodontidae, from the Mediterranean Coast of Egypt. Egyptian J. of Aquatic Research 31(I):
- Erzini, K. (1991). A compilation of data on Variability in length-age in marine fishes". Working paper 77. Fisheries stock Ssimnort
- Faltas, S. N. (1993). Studies on the fishery biology of lizard fish (Family: Synodontidae) in the Egyptian Mediterranean waters. Ph.D. Thesis, Fac. Sci., Alexandria University, Egypt.
- Federizon, R. (1993). Using vital statistics and survey catch composition data for tropical multispecies fish stock assessment: application to the demersal resources of the central Philippines. Alfred-Wegener-Institut für Polar- und Meeresforschung, Bremerhaven. Ph.D. dissertation, 201pp.
- Gayanilo, J.; Sparre, F. C. P. and Pauly, D. (1997). The FAO-ICLARM Stock Assessment Tools (FISAT) "FAO Computerized Information Series (Fisheries).No.8.Rome, FAO.
- Golani, D. (1993). The biology of the Red Sea migrant, *Saurida undosquamis* in the Mediterranean and comparison with the indigenous confamilial *Synodus saurus* (Teleostei: Synodontidae). Hydrobiologia 271(2):109-117.

- Gulland, J. A. (1971). *The Fish Resources of the Ocean*. West Byfleet, Surrey, Fishing News (Books), Ltd., for FAO; 255pp.
- Gulland, J. A. and Holt S. I. (1959). Estimation of growth parameters for data at unequal time intervals .J. Cons. Perm. Int. Explor Mer,25 (1):47-49.
- Ismen, A. (2003). Maturity and fecundity of lizard fish (*Saurida undosquamis* Richardson, 1848) in Iskenderun Bay (Eastern Mediterranean) . Turk. J. Zool., 27: 231-238.
- Latif, A. A. and Shenouda, T. S. (1973). Studies on *Saurida undosquamis* (Richardson) from the Gulf of Suez. Monthly peculiarities of gonads. Bull. Inst. Oceanogr. Fish. Egypt., 3: 295-335.
- Pauly, D. (1983a). Length-converted catch curves. A powerful tool for fisheries research in the tropics. Part 1. ICLARM Fishbyte, 1(2): 9-13.
- Pauly, D. (1983b). A selection of simple methods for the assessment of tropical fish stocks. FAO Fish. Circ. (729): 54 pp.
- Pauly, D. (1984a). Length-converted catch curves. A powerful tool for fisheries research in the tropics. (part II) ". ICLARM Fishbyte, 2 (1): 17-19. .
- Pauly, D. (1984b). Length-converted catch curves. A powerful tool for fisheries research in the tropics. (III: conclusion) ". ICLARM Fishbyte, 2 (3): 9-10.
- Pauly, D., and Munro, J. L. (1984). Once more on the comparison of growth in fish and invertebrates. ICLARM Fishbyte, 2 (1): 21
- Pauly, D. and Soriano, M. L. (1986). Some practical extensions to Beverton and Holt's relative yield-per-recruit model, p. 491-496. In J. L. Maclean, L. B. Dizon and L. V. Hosillo (eds.). *The First Asian Fisheries Forum*. Asian Fisheries Society, Manila , Philippines.
- Pitt T. K. (1970). Distribution, abundance and spawning of yellowtail flounder, *Limanda ferruginea*, in the New Foundland area of the North West Atlantic. J. Fish. Res. Bd. Canada, 27(12): 2261-2271.
- Ramadan, A. M. (1995). Reproduction studies on Lizard fish, *Saurida undosquamis*. M. Sc., Thesis, Faculty of Science. Suez Canal University.

- Rao K.V. S. (1983). Maturation and spawning of lizard fishes (*Saurida spp.*) from North Western part of Bay of Bengal. *Indian J. Fish*, 30(1): 27-45.
- Sanders, M. J.; Kedidi, S. M. and Hegazy, M.R. (1984). Stock assessment for the brushtooth lizardfish (*Saurida undosquamis*) caught by trawl in the Gulf of suez. Project for the Development of Fisheries in the areas of the Red Sea and Gulf of Aden", FAO/UNDP RAB/83/023/05. Cairo. 28 pp. (mimeo)
- Siripakhavanich, S. (1990). A study on population dynamics of the brushtooth lizard fish, *Saurida undosquamis* (Richardson) off the upper western coast in the Gulf of Thailand. " Faculty of Fisheries, Kasetsart University, Thailand., 77 pp.
- Sousa, M. I. (1992). Seasonal growth of five commercially important fishes at Sofala Bank, Mozambique". *Rev. Invest. Pesq. (Maputo)*, 27: 79-97.