

Monitoring of 17 β -Estradiol Residues in the Suez Canal Region

Nagwa Elnwishy¹; Amro Hanora²; Martin Hedström³; Raafat Afifi⁴;
Bo Mattiasson³ and Helmy Omran⁵

- 1- Biotechnology Research Center (BRC), Suez Canal University, (New Campus), Ismailia 41522, Egypt.
 - 2- Department of Microbiology and Immunology, Faculty of Pharmacy, Suez Canal University, Ismailia 41522, Egypt.
 - 3- Department of Biotechnology, center of chemistry & chemical engineering, Lund university, Lund 22100, Sweden.
 - 4- Department of Marine Biology, Faculty of Science, Suez Canal University, Ismailia 41522,, Egypt.
 - 5- Department of Food Technology, faculty of Agriculture, Suez Canal University, Ismailia 41522,, Egypt.
- E-mail: nwishy@yahoo.com

ABSTRACT

This study was initiated to provide the first record of monitoring of 17 β -estradiol (E2) residues in some Egyptian aquatic ecosystems. Samples of water were collected from three water bodies located in the Suez Canal region. Samples were extracted, filtered and examined by HPLC on a C₁₈ column using Florescence detectors.

Results provided evidence of the presence of estradiol in the studied area. Interestingly, marine lakes contained significant levels of 17 β -estradiol (P <0.05). Lower levels were also detected in the rivulet streams supplied by River Nile.

Detection of estradiol in the aquatic ecosystems of the Suez Canal region grabs the attention towards the heavy reliance on some esterogenic medicinal products in the area, and the eventual effect on the aquatic systems including biodiversity of a variety of organisms. Therefore, it is recommended to enlarge the detection scanning of estradiol in other Egyptian areas.

Key words: Estradiol, Temsah Lake, Suez Canal region

INTRODUCTION

There is a global concern about the presence of the estrogenic residues in the aquatic ecosystems as one of the dangerous pharmaceutical residues; this is due to its impact on marine life ecosystems and on human health (Seifert *et al.*, 2003). The source of these estrogenic residues is industrial wastes and medicines (Nolan *et al.*, 2001), and as additives in animal feed (Bruce, 2005; Hailing *et al.*, 2002). Even though that these compounds are found in small amounts in the environment (Hensen *et al.*, 2002), but the effect of these traces is remarkable on marine animals and consequently on humans.

It was evidenced that estrogenic residues were detected in some lakes, rivers and ground waters in many places around Europe (Burton *et al.*, 2002). It was also reported that it caused severe damage to the marine system balance and on human welfare.

17 β estradiol (E2) is one of these compounds which belong to (EDCs) and have been of important concern since the 1930s (Cook *et al.*, 1934; Tawfic, 2006). 17 β Estradiol, as well as other steroids, are excreted from the human body and live

stocks in high quantities (Narender and Cindy, 2009); so when they are discharged into marine environments as waste waters or sewage, they occur with concentration levels in the range of nanograms /L (Koh *et al.*, 2008). Concentrations less than 1 ng/L in the aquatic environment can cause infertility and weaken estrogenic activity, decrease reproductivity of female fish (Burton and Wells, 2002), change behavioral habits which may change reproductive physiology of fish (Denslow and Sepúlveda, 2007), and sex reverse in some fish turning males to females (Hansen *et al.*, 1998).

Since contraceptives have been widely used in the last 50 years in many countries, 17 β estradiol, and its derivatives, were detected in aquatic environments in different industrial countries, even where high precautions of water and environmental safety are considered. They were detected in British raw domestic sewage discharged into rivers (Desbrow *et al.*, 1998; Rujiralai *et al.*, 2011) and waste water in South Korea (Ra *et al.*, 2011), China (Liu *et al.*, 2011; Lu *et al.*, 2011; Ra *et al.*, 2011; Zhou *et al.*, 2011), The Netherlands (Belfroid *et al.*, 2006), Italy (Pojana *et al.*, 2007), and Germany (Körner *et al.*, 2001; Hintemann *et al.*, 2006). In addition, it was detected in prawn and fish (Jiang *et al.*, 2009), Japanese Spanish mackerel (*Scomberomorus niphonius*) and mollusks (Zou *et al.*, 2007), in France, it was also detected in shellfish (Lagadic *et al.*, 2007). Therefore, the aim of this project was to investigate and locate 17 β estradiol residues in Suez canal region as a first step to scan other Egyptian aquatic ecosystems.

MATERIALS AND METHODS

Study areas:

The study was carried out in the Suez Canal region (Fig. 1). Samples were collected from Bahr Elbakar drainage, Ismailia stream, Abo shehata., and from two parts of Temsah Lake which consists of two different ecosystems together, a fresh water part supplied from the River Nile, and a marine water part supplied by the Suez canal. In addition to Temsah Lake, samples were collected from Manzala lake. A volume of 100 ml were collected at 50 cm depth, filtered, then stored frozen until they were thawed as a pre-step to analysis.

Quantitative verification of target Residues:

Extraction of estradiol

A volume of 100 ml of each sample was freeze-dried. (Hetovac). The remaining dried material was re-dissolved in 6 ml of n-hexane (Merck, Lichrosolv) and then shaken vigorously for 30 minutes on a shaker.

Clean up of the samples:

Molecular Imprinted Polymer (MIP) was prepared according to Le Noir *et al.*, (2007), and packed in a glass columns. Just before clean up, the MIP columns were conditioned with 5 ml of methanol: acetic acid (2:1), at a flow rate of 1ml/ 10 min. Hexane extract from each sample was applied to MIP column at a flow rate of 1ml/ min. Finally, the column was eluted with 10 ml of a mixture of methanol: acetic acid (2:1) at a flow rate of 0.5 ml /minuet (Le Noir *et al.*, 2007).

The collected samples were concentrated to 500 μ l by evaporation under nitrogen gas stream. The final samples were filtered through 0.45 μ m filters and transferred to 2 ml vials and estradiol concentrations were determined by HPLC system using C₁₈ column and and florescence detectors.




Fig. 1: location of sampling sites in Suez Canal region

Detection by HPLC- fluorescence analysis

A series of 17 β -estradiol (98% purity, Sigma Aldrich) dilutions (2×10^6 ng/ml-0.0001 ng/ml) were analyzed to construct a standard curve, and the cleaned up samples were subjected to HPLC- fluorescence for detection under the same conditions. Results were evaluated using the standard curve to determine the concentrations in the tested samples.

The analysis was made using an Auto-sampler G13129, and the separation was carried out on Supelcosil C₁₈ column (150 x 4.60 mm, 5 μ m). Elution was performed with gradient elution using acetonitrile 10% in ultra pure water up to 100% acetonitrile (Merck, Lichrosolv-for chromatography) and water at flow rate of 0.8 ml/min (Lopez de Alda *et al.*, 2002). Estradiol was detected using fluorescence detector G 1321A at excitation at 420 nm after emission at 330 nm. Detection was possible in the range 1.0 mg/ml -0.01 ng/ml.

Statistical analysis:

The statistical analysis was performed by applying ANOVA test to the results to compare differences between sites. The correlation between samples was performed using IBM SPSS Version19, 2010.

RESULTS

Estradiol was spiked up at minute 23.68 in the standard curve as well as in the contaminated samples as shown in Figure 2. Estradiol was detected in all tested samples as summarized in Table (1) and illustrated in Fig (3). The highest level of estradiol ($1029 \pm 0.1 \mu\text{g/L}$) was found in Bahr Elbakar samples. Also, it was detected in both marine and fresh water parts of Tamsah Lake; (925 ± 0.1 and $462 \pm 0.1 \mu\text{g/L}$)

respectively). Lower concentrations were detected in Ismailia stream, El Salam canal, and Abo Shehata (556 ± 0.1 , 552 ± 0.1 , and 284 ± 0.1 $\mu\text{g/L}$) respectively.

Table 1: Detected concentrations of 17 β -estradiol (\pm St. Div.)

Location	Sample	$\mu\text{g/L}$	St. Div.
Bahr El Bakar	w9	1029	0.1
Manzala Lake	w10	684	0.1
Salam canal	w12	552	0.1
Temsah lake "Marine"	w13	925	0.1
Temsah lake "fresh"	w14	462	0.1
Ismailia stream	w15	556	0.1
Abo Shehata	w16	284	0.1




Fig. 2: HPLC analysis showing the standard of 17 β estradiol (a) and its detection in Bahr elbakar sample (b) at retention time 23.684.




Fig. 3: Detected concentration in of 17 β estradiol in the tested samples

DISCUSSION

Bahr Elbakar drain had the highest level of estradiol ($1029 \pm 0.1 \mu\text{g/L}$) as this drain is the end point stream of Cairo and Delta discharges. This high detected concentration must be because the drain receives untreated and somehow primarily treated waste water from east Cairo moving east to Manzala lake (Taha *et al.*, 2004). In addition, the industrial activities, waste water discharged from Belbeis drain to Bahr El-Bakar drain, as well as domestic discharge from the country side around Bahr El-Bakar drain, all these sources may have resulted in having such high concentrations of estradiol.

E2 detection in the marine and the fresh water parts of Temsah Lake; (925 ± 0.1 and 462 ± 0.1 respectively) is interesting because that lake is unique as it receives high saline water supply from the Suez Canal moving from south (Red sea) to north (Mediterranean sea), and receives freshwater supply as a mixture coming from the Nile and other sub-Nile streams (El-Serehy and Sleight, 1992). The lake is an important source of feeding for many living species; most of the aquatic migrating birds stop around Temsah lake in their migratory trip from Europe to Africa in winter and vice versa in summer. Local birds also live around the lake and its surrounding areas, where they find their feeding resources of fish and crustaceans. People in this area depend on fish as their main source of food. The annual fish production from Temsah lake and the bitter lakes is 4557 tons of the total fish production in Egypt which is estimated with 387398 tons according to GAFRD (2009). However, Temsah lake seems to be in need of proper attention and conservation. The lake has been suffering from different sources of pollution for decades endangering the environment, marine life and human welfare (El-Sherif *et al.* 2009).

The concentration of estradiol detected in Manzala lake was $684 \pm 0.1 \mu\text{g/L}$. This lake is also known to be the end point of different discharges of the surrounding area though it contributes with 48023 tons to the total Egyptian fish production (GAFRD, 2009). In addition, the lake is the final destination of 3 billion m^3 of waste water discharge coming from Bahr Elbakar drain every year (EU-SMAP, 2007). Thus, the water quality in Manzala Lake is affected by the water quality received from Bahr Elbakar. However, this concentration is in one way or another below the expected concentrations, taking into consideration the huge amount of polluted discharges in the lake. But this may be due to the dilution effect by the water flow from Suez Canal towards the Mediterranean sea, or most likely due to the absence of medical industrial discharges or animal farms' drainage which usually contains high amounts of

estrogens incorporated in the animal feed as additives (Bruce, 2005; Hailing *et al.* 2002), unlike the case in Tamsah lake which is surrounded by agricultural farms, and few animal production activities by the farmers.

Furthermore, the lower concentrations of estradiol detected in El Salam canal, Ismailia stream and Abo Shehata (552 ± 0.1 , 556 ± 0.1 and 284 ± 0.1 $\mu\text{g/L}$) respectively is most likely because they are bronchial streams and canals where the dilution factor through hundreds of kilometers of water flow away from the main sources would have a considerable effect. For instance, El Salam canal, which is an artificial canal established to supply eastern parts of the Suez canal region with irrigation water, is 262 Km^2 long. It is supplied by a mixture of 2.11 billion m^3/year of the Nile fresh water from the Domietta branch, 1.905 billion m^3/year of the drainage water from Bahr Hadous and 0.435 billion m^3/year of El Serw drainage water (Hafez, 2005). Thus, the ratio of Nile water to drainage water is approximately 1:1. However, the water is used in fish farming and animal feeding as additional farm activities (Elnwishy *et al.*, 2008) besides being used in irrigation for reclamation of east Suez canal which is the major purpose for establishing the canal (Hafez, 2005). The mixture of the sewage water and the farm's discharges may have resulted in some harmful biological effects on fish and animal reproduction in the area.

Also, the concentrations in Ismailia stream (556 ± 0.1 $\mu\text{g/L}$) which is supplied by Nile fresh water from north Cairo to Ismailia were higher than that detected in the fresh water part of Tamsah lake (462 ± 0.1 $\mu\text{g/L}$) though they are supplied by the Nile and closely located branches. But this may be due to a higher load of discharges in the lake, while on the other hand, the dilution factor to the discharges in the stream from Cairo to Ismailia slightly reduce the concentration.

It was observed that the tested samples from the marine water contained higher concentrations of estradiol than the samples from fresh and brackish waters. The marine part of Tamsah lake had an approximately 50 % higher contamination level of estradiol than the fresh water part of the same lake, and both Tamsah and Manzala lakes showed higher concentrations of estradiol than other samples. It is worth to mention that the salinity in both lakes are ~ 40 ‰ (El-Serehy and Sleigh, 1992). Different factors may also be involved in such higher level in marine water than fresh water; such as faster degradation of estradiol in the anaerobic freshwater sediments than in the anaerobic marine sediments (Lopez de Alda *et al.*, 2002; Tyler *et al.*, 2005; Czajka and Londry, 2006; Christoph and Juliane, 2009), salinity assistance to estradiol to become more resistant to degradation, or the inability and sometimes total failing of marine microorganisms to be involved in biodegradation of estradiol are all possible factors open for upcoming investigations.

CONCLUSION

It was concluded that the Suez canal region is contaminated with 17 β -estradiol residues. Tamsah and Manzala lakes showed relatively high concentrations. Though these levels are not extremely high but, they may result in unhealthy effects to human consumption of sea food due to accumulation factor.

However, this research provides the first data about the presence and the levels of estradiol in Egypt. Thus, further investigations are recommended in the future in other regions in Egypt as well as to adopting new techniques for the removal of 17 β -estradiol in order to maintain marine life ecosystem in the tested locations.

ACKNOWLEDGMENT

The authors would like to acknowledge the Erasmus Mundus External Co-operation Window (EMECW), and the Swedish Research Link program for financial support, and the Biotechnology Research Center in Suez Canal University, for providing laboratorial facilities.

REFERENCES

- Belfroid, A.C.; Schrap, S.M. and de Voogt, P. (2006). Occurrence of estrogenic hormones, bisphenol-A and phthalates in the aquatic environment of The Netherlands, in: Vethaak, A.D. *et al.* (Ed.) Estrogens and xenoestrogens in the aquatic environment: an integrated approach for field monitoring and effect assessment, pp. 53-75.
- Burton, J.L. and Wells, M. (2002). The effect of phytoestrogens on the female genital tract. *Journal of Clinical Pathology* 55:401-407.
- Bruce S. (2005): Water Resource Impacts from Medicines and Other Biologically Active Substances, WQ-1278, North Dakota State University, Fargo. <http://www.ag.ndsu.edu/pubs/h2oqual/watgrnd/wq1278.pdf>
- Christoph, M. and Juliane, H. (2009). Microbial Degradation of Steroid Hormones in the environment and Technical Systems. Lecture, http://www.ibp.ethz.ch/research/aquaticchemistry/teaching/archive_past_lectures/term_paper_08_09/HS08_CHRISTOPH_MOSCHET_rev_termpaper.pdf.
- Cook, J.W.; Dodds, E.C.; Hewett, C.L. and Lawson, W. (1934). The oestrogenic activity of some condensed-ring compounds in relation to their biological activities. . *Proc. Roy. Soc. Lond.*, 114, 272-286. Cited in "Koh, Y. K. K.; Chiu, T. Y.; Boobis, A.; Cartmell, E.; Scrimshaw, M. D. and Lester, J. N. (2008). "Treatment and removal strategies from wastewater" *Environmental Technology* 2029(2003): 2245-2267.
- Czajka, C.P. and Londry, K.L. (2006). Anaerobic biotransformation of estrogens. *Science of The Total Environment* 367: 932-941.
- Denslow, N. and Sepúlveda, M. (2007). Ecotoxicological effects of endocrine disrupting compounds on fish reproduction. *The Fish Oocyte*. In: Babin, P.J., Cerdà, J., Lubzens, E. (Eds.). Springer Netherlands, pp. 255-322.
- Desbrow, C.; Routledge, E.J.; Brighty, G.C.; Sumpter, J.P. and Waldock, M. (1998). Identification of Estrogenic Chemicals in STW Effluent. 1. Chemical Fractionation and *in vitro* Biological Screening. *Environmental Science & Technology* 32: 1549-1558.
- El-Serehy, H.A.H. and Sleigh, M.A. (1992). Physico-chemical conditions for plankton in Lake Timsah, a saline lake on the Suez Canal. *Estuarine, Coastal and Shelf Science* 34: 127-139.
- El-Sherif, M.S.; Ahmed, M.T.; El-Danasoury, M.A. and Elnwishy, N. (2009). Effects of pollutants on some aquatic organisms in Timsah Lake in Egypt. *Journal of Fisheries & Aquatic Science*. 4(3):150-160
- Elnwishy, N. Saleh, M. and Zalat, S. (2008) Combating desertification through fish farming. *The Future of Drylands 2006: International Scientific Conference on Desertification and Dry Lands Research*, UNESCO publishing, pp. 507-518
- EU-SMAP, European Union Short and Medium-Term Priority Environmental Action programme (2007). Action 6: The Execution of Specific Assessments in the Coastal Areas of Port Said. Irrigation Advisory Service – Ministry of Water Resources and Irrigation & Resources Technology Bureau (RTB). Contract n. MED/2005/112-172

- GAFRD, General Authority of Fish Resources Development (2009). Report of annual production of fish in the Egypt for 2009. General authority of fish resources development.
- Hafez A. (2005). Investigation of El-Salam Canal Project In Northern Sinai, Egypt Phase-I: Environmental Baseline, Soil And Water Quality Studies. Ninth International Water Technology Conference, IWTC9 2005, Sharm El-Sheikh, Egypt. pp. 953-970.
- Halling-Sorensen, B.; Nielsen, S. and Jensen, J. (2002). Environmental assessment of veterinary medicinal products in Denmark. Environ. Proj.No. 659 (2002). Danish Environmental Protection Agency.WQ-1278 Water Resource Impacts from Medicines and Other Biologically Active Substances Stuer-Lauridsen, F., M. Birkved,
- Hansen, P.; Dizer, H.; Hock, B.; Marx, A.; Sherry, J. and McMaster, Blaise, C. (1998): Vitellogenin- a biomarker for endocrine disruptors. Trends in Analytical Chemistry 17: 448-451.
- Hintemann, T.; Schneider, C.; Schöler, H.F. and Schneider, R.J. (2006). Field study using two immunoassays for the determination of estradiol and ethinylestradiol in the aquatic environment. Water Research, 40:2287-2294.
- Jiang, T.; Zhao, L.; Chu, B.; Feng, Q.; Yan, W. and Lin, J.-M. (2009). Molecularly imprinted solid-phase extraction for the selective determination of 17[beta]-estradiol in fishery samples with high performance liquid chromatography. Talanta 78: 442-447.
- Koh, Y.K.K.; Chiu, T.Y.; Boobis, A.; Cartmell, E.; Scrimshaw, M.D. and Lester, J.N. (2008). Treatment And Removal Strategies For Estrogens From Wastewater. Environmental Technology 29:245-267.
- Körner, W.; Bolz, U.; Triebkorn, R., Schwaiger, J., Negele, R.-D., Marx, A., and Hagnemaier, H., (2001). Steroid analysis and xenosteroid potentials in two small streams in southwest Germany. Journal of Aquatic Ecosystem Stress and Recovery (Formerly Journal of Aquatic Ecosystem Health) 8: 215-229.
- Lagadic, L., Coutellec, M.-A., and Caquet, T.,(2007). Endocrine disruption in aquatic pulmonate molluscs: few evidences, many challenges. Ecotoxicology 16: 45-59.
- Le Noir, M.; Lepeuple, A.-S.; Guieysse, B. and Mattiasson, B. (2007). Selective removal of 17[beta]-estradiol at trace concentration using a molecularly imprinted polymer. Water Research 41: 2825-2831.
- Liu, S.; Ying, G.-G.; Zhao, J.-L.; Chen, F.; Yang, B.; Zhou, L.-J. and Lai, H.-j. (2011). Trace analysis of 28 steroids in surface water, wastewater and sludge samples by rapid resolution liquid chromatography-electrospray ionization tandem mass spectrometry. Journal of Chromatography 1218 (A): 1367-1378.
- Lopez de Alda, M.J.; Gil, A.; Paz, E. and Barcelo, D.(2002). Occurrence and analysis of estrogens and progestogens in river sediments by liquid chromatography-electrospray-mass spectrometry. Analyst 127|: 1299-1304.
- Lu, G.; Yan, Z.; Wang, Y. and Chen, W. (2011). Assessment of estrogenic contamination and biological effects in Lake Taihu.
- Narender, K.N. and Cindy, L.M. (2009). Water Quality Guidelines for Pharmaceutically-active-Compounds (PhACs): 17 α -ethinylestradiol (EE2). Report by Ministry of Environment province of British Columbia, <http://www.env.gov.bc.ca/wat/wq/BCguidelines/PhACs-EE2/PhACs-EE2-tech.pdf>.
- Nolan, M.; Jobling, S; Brighty, G. and Sumpter, J. (2001): A histological description of intersexuality in the roach. J. Fish Biol. 58:160–176.
- Pojana, G.; Gomiero, A.; Jonkers, N. and Marcomini, A. (2007). Natural and synthetic endocrine disrupting compounds (EDCs) in water, sediment and biota of a coastal lagoon. Environment International 33: 929-936.

- Ra, J.-S.; Lee, S.-H.; Lee, J.; Kim, H.Y.; Lim, B.J.; Kim, S.H. and Kim, S.D. (2011). Occurrence of estrogenic chemicals in South Korean surface waters and municipal wastewaters. *Journal of Environmental Monitoring* 13: 101-109.
- Rujiralai, T.; Bull, I.D.; Llewellyn, N. and Evershed, R.P. (2011). In situ polar organic chemical integrative sampling (POCIS) of steroidal estrogens in sewage treatment works discharge and river water. *Journal of Environmental Monitoring* 13: 1427-1434.
- Seifert, M.; Li, W.; Alberti, M.; Ausch, U. and Hock B. (2003): biomonitoring: Integration of biological endpoints into chemical monitoring. *ppl. Chem.*, 7(11-12) pp.2451 –2459.
- Taha A. A.; El-Mahmoudi A. S. and El-Haddad I. M (2004). Pollution Sources and Related Environmental Impacts in the New Communities Southeast Nile Delta, Egypt. *Emirates Journal for Engineering Research*, 9 (1): 35-49
- Tawfic, M.(2006). Persistent Organic Pollutants in Egypt- An Overview. *Soil and Water Pollution Monitoring, Protection and Remediation*. In: Twardowska, I., Allen, H.E., Häggblom, M.M., Stefaniak, S. (Eds.). Springer Netherlands, pp. 25-38.
- Tyler, C.R.; Spary, C.; Gibson, R.; Santos, E.M.; Shears, J. and Hill, E.M. (2005). Accounting for differences in estrogenic responses in rainbow trout (*Oncorhynchus mykiss* : Salmonidae) and roach (*Rutilus rutilus* : Cyprinidae) exposed to effluents from wastewater treatment works. *Environmental Science & Technology* 39:2599-2607.
- Zhou, Y.; Zha, J.; Xu, Y.; Lei, B. and Wang, Z. (2011). Occurrences of six steroid estrogens from different effluents in Beijing, China. *Environmental Monitoring and Assessment*, pp.1-11.

ARABIC SUMMARY

رصد بقايا مركب 17 - بيتا استراديول في منطقة قناة السويس

- نجوى النويشى¹، عمرو حنورة²، مارتن هيدسترومب³، رأفت عفيفي⁴ بو ماتياسون³، حلمي عمران⁵
 1- مركز بحوث التقنية الحيوية، جامعة قناة السويس، 41522 الإسماعيلية، مصر.
 2- قسم الميكروبيولوجيا والمناعة، كلية الصيدلة، جامعة قناة السويس، الإسماعيلية 41522، مصر.
 3- قسم التكنولوجيا الحيوية، ومركز الهندسة الكيميائية والمواد الكيميائية، جامعة لوند، لوند 22100، السويد.
 4- قسم علم الأحياء البحرية، كلية العلوم، جامعة قناة السويس، الإسماعيلية 41522، مصر.
 5- قسم تكنولوجيا الأغذية، كلية الزراعة، جامعة قناة السويس، الإسماعيلية 41522، مصر.
 * بريد إلكتروني nwishy@yahoo.com

هذه الدراسة تسجل اول رصد لمتبقيات مركب 17 بيتا استراديول في بعض النظم الايكولوجية المائية المصرية. حيث تم جمع عينات من المسطحات المائية في منطقة قناة السويس. وتم استخلاص العينات وفحصها باستخدام جهاز HPLC على عمود C18 باستخدام أجهزة الكشف عن التلوين. وقد قدمت النتائج أدلة على وجود الاستراديول في المنطقة التي شملتها الدراسة. ومن المثير للاهتمام، وجود مستويات كبيرة من 17- β استراديول ($P < 0.05$) في البحيرات المالحة عنها في المياه العذبة. ورصدت أدنى المستويات في تيارات المصارف المائية المتفرعة من نهر النيل. وتبين وجود الاستراديول في النظم الايكولوجية المائية في منطقة قناة السويس يجذب الانتباه نحو خطورة الاعتماد الكبير على بعض المنتجات الطبية الستيرويدية في المنطقة، والتأثير النهائي لوجودها على النظم المائية بما في ذلك التنوع البيولوجي من مجموعة متنوعة من الكائنات الحية. ولذلك فمن المستحسن توسيع نطاق عملية الكشف عن الاستراديول في مسطحات مائية اخرى في جمهورية مصر العربية.