

**GROWTH PERFORMANCE OF FISH REARED UNDER
DIFFERENT DENSITIES IN SEMI-INTENSIVE
AND EXTENSIVE EARTHEN PONDS**

**Ibrahim M. Shaker Abed El-Fattah¹, and
Mohamed M. Abdel-Aal.²**

1- Limnology Department and

2- Aquaculture Department Central Laboratory for Aquaculture
Research, Abbassa, Sharkia, Egypt.

Key words: polyculture, intensive, extensive, water quality, plankton,
growth performance, tilapia, mullet, carp, catfish.

ABSTRACT

Effect of stocking density on fish performance has been investigated in earthen ponds over five months of growing period in a polyculture system including Nile tilapia, common carp, silver carp, mullet and African catfish. Experimental ponds have been allocated to various stocking rates of 1, 1.25, 1.5, 1.75, 10 and 14 fish/m³. The treatments of 10 and 14 fish/m³ represented the semi-intensive system, while lower densities represented the extensive culture system. Fish in all treatments received 25% crude protein pelleted fish feed at a rate of 3% of body weight that was adjusted throughout the growing period. Feed was offered twice/day for five days/week. Water quality has been monitored monthly. The physico - chemical analysis showed two distinct sets of values for the extensive and the semi-intensive stocking densities concurred with for the main water quality parameters, whereas the values of pH, E.C., salinity, alkalinity, hardness, total phosphorus and orthophosphate were significantly higher in extensive system than the semi-intensive one, while values of D.O, SD, NH₃, NO₂, NO₃ were higher in the semi-intensive system compared to the extensive system. Plankton communities expressed as Chlorophyll "a" was significantly higher in the extensive ponds. Average production was significantly higher in semi-intensive ponds compared to the extensive ponds. However, for the main fish species (Nile tilapia), the average weight as well as the daily gain did not vary among treatments, while there were some differences with regard to average weight for other species. There was no difference concerning the condition factor in all treatments for any of the tested species. The

economic analysis revealed increases in variable costs as well as higher net returns as the stocking density increased. However, the rate of return to capital percentage has decreased with the increase of the stocking density. The overall analysis suggested the semi-intensive system to be the most profitable system.

INTRODUCTION

The polyculture as practiced in aquaculture via stocking of fish species of different food habits proved to be an important management tool in utilizing efficiently the natural food resources in fish pond. Synergistic interactions among fish species are manifested by higher growth and yield in polyculture than in monoculture (Karplus *et al.* 1996). The bases for these interactions are the increase of available food resources and the improvement of environmental conditions (Milstein, 1992).

Polyculture was first practiced in China more than a thousand years ago. It was extremely extensive, requiring little management, stocking several species at low density, often without application of feeds, and producing relatively low yield at low production costs (Lin, 1982). In semi-intensive farming systems, usually natural food is stimulated through the application of organic manures or chemical fertilizers, while feeds are supplied to supplement the available natural food. Aeration devices and automatic feeders have been incorporated into such system whenever possible. By definition, fish are stocked in the semi-intensive system at higher densities compared to that in the extensive ones, and thus, production costs as well as yields are expected to be much higher. In these systems, the supplemented organic wastes and feeds are utilized directly by the fish and also act as fertilizers for the heterotrophic and autotrophic food web. Under manuring and supplemental feeding, natural food still represents an important component of the overall food utilized by fish. The autotrophic path constitutes not only the food source of plankton filter feeding fish, like silver carp, but also provides 60-80% and 50% of tilapia and carp requirements, respectively. The fish growth can even be attributed to natural food developed on the pond bottom (karplus, 1996).

Semi-intensive polyculture and monoculture systems are widely used in fish culture in Egypt, practiced in shallow earthen ponds and in deep dual-purpose irrigation reservoirs. The main species cultured in commercial ponds are Nile tilapia, *Oreochromis niloticus*, common carp

Cyprinus carpio, silver carp, *Hypophthalmichthys molitrix*, striped mullet, *Mugil cephalus* and grey mullet, *Liza ramada*. These species are stocked in variety of combinations. The outcomes of such system as fish biomass vary widely.

Recently, along with the intensification trend in Egyptian aquaculture, the requirement for high quality feed has increased. Such higher production costs as well as the high investments have called for better management practices. The contribution of feed costs in fish production increases along with the level of intensification, reaching the highest component in farm operation (Islam, 2002). From economic point of view, such practices may not suite the small-scale farmers with little resources whether in the tropics and to some extent in some parts of Egypt. This is due to financial constraints which limit the ability of those farmers from accessing the commercially produced pelleted feeds. Thus, in order for those farms to utilize easy to find feed at low cost, alternative management strategies are required.

Fish stocking density is a key factor in determining the management of fish pond and so, the production. It affects the amounts of natural food available per fish, the level of supplemented feeding required (Hepher, 1988), and thus, the intensity of inter- and intra-specific food competition. With increasing stocking density, competition for food among fish increases. The optimal combined stocking densities of several species of fish cultured together should be carefully analyzed in order to optimize both fish performance (growth and yield) and economic profitability of the combined system. Therefore, the objectives of the present study are to evaluate fish production and growth performance as well as the grow-out environments under different stocking densities of fish cultured. Economical evaluation for farming systems tested was a main element in the study. Finally, the study tackled the issue of the use of drainage water in Egyptian fish farming.

MATERIALS AND METHODS

The present study was conducted in earthen ponds; including six ponds of 4,200 m² surface area - each and twelve ponds of 54,600 m² surface area each. Ponds were located at Manzala fish farm, Manzala, Egypt. The first six ponds had 1.5 m as an average water depth, while the twelve ponds had an average water depth of 1.25 m. Aeration devices were used in the six ponds (2 Impeller Paddle wheel Aerator-AR-A232-2 h), while, no aeration was practiced in the twelve ponds. Fish were

stocked on in May and harvested in 20 October 2005 after about 150 days of growth period. Each treatment was conducted in three replicates.

The present study consisted of two parts, which were conducted in completely randomized design. The first group was used to evaluate the fish stocking density under the semi-intensive system at a rate of 10 and 14 fish/m³. The second part of the study was targeted to evaluate the lower stocking density used in the extensive system, where the rate of 1, 1.2, 1.5 and 1.7 fish/m³ have been used as shown in Table 1. The two systems had been stocked with the same fish species, namely, Nile tilapia, (*Oreochromis niloticus*); common carp, (*Cyprinus carpio*), silver carp, (*Hypophthalmichthys molitrix*), mullet, (*Mugil cephalus*) and catfish, (*Clarias gariepinus*). Commercial feed pellets (25% crude protein) was used throughout the experiment and was provided twice a day; five days a week at a rate of 3% of body weight and automatically and manually distributed for the semi-intensive and extensive fish ponds, respectively. All experimental ponds were fertilized using poultry liter (150 kg/Fadden¹). Thereafter, ponds were filled with water for no longer than two weeks before fish stocking. Fish sampling for all species was carried out monthly and the amount of feed was adjusted accordingly.

Water sampling was carried out for several parameters of concern to aquaculture. Physico-chemical parameters were monitored monthly. Temperature and dissolved oxygen were determined directly by a portable digital oxygen meter (YSI model 58, USA), pH was measured using a digital pH meter (Accumet 340) and transparency by a Secchi disc. Water electrical conductivity and salinity were determined by conductivity meter (Orion 630), Nitrate, nitrite and free ammonia were determined using a HACH water analysis kits (DR 2000, USA), while total phosphorus, orthophosphate, total alkalinity, total hardness, chlorophyll "a" were determined according to standard methods (APHA, 2000) 100ml water sample was filtered through 0.45 M Millipore filter and chlorophyll a was extracted in 5 ml of 90% acetone and grinded by tissue grinder, kept for 24h at 5°C. The sample was then centrifuged and measured the absorbance of acetone, chlorophyll a concentration was calculated using the equation:

$$\text{Chlorophyll a in } \mu\text{g} = 11.9(A_{665} - A_{750}) V/L \times 1000/s$$

Where: A₆₆₅= the absorbance at 665A, A₇₅₀ the absorbance at 750, V= the acetone extract in ml, L= the length path in the spectrophotometer in cm, S= the volume in ml of sample filtered.

¹ Fadden = 4200 m²

Qualitative and quantitative estimates of phytoplankton and zooplankton were also recorded monthly. The analyses of heavy metals in water and different fish species were determined using atomic absorption (Thermo 6500,) with graphite furnace according to APHA (2000). At the end of the experiment, fish were harvested, counted and weighed. The growth parameters were calculated as follows:

Daily gain (DG) = $(Wt_2 - Wt_1) / T$; Specific growth rate (SGR) = $(\ln Wt_2 - \ln Wt_1) \times 100 / T$; where Wt_1 is the initial weight in grams, Wt_2 is the second weight in grams, and T is the period in days Condition factor (K) = $\text{Body weight} / \text{Total length}^3 \times 100$.

Statistical analysis was performed using the analysis of variance (ANOVA). Duncan's Multiple Range Test Duncan (1955) was used to determine the significant differences between means at $P < 0.05$. Standard errors of treatment means were also estimated. All statistical evaluations were carried out using Statistical Analysis Systems (SAS) program (SAS, 2000) according to Steel and Torrie (1960).

RESULTS AND DISCUSSION

The results of physico-chemical parameters and the selected heavy metals in water under different experimental treatments are shown in Tables (2 and 3). The pH values in pond water of the treatments ranged from 8.45 to 9.6. This variation could be explained by the photosynthetic uptake of CO_2 and bicarbonate that substituted hydroxyl ions. The pH values in extensive ponds were not significantly different. These results indicate that the increase of fish density from 1.0 to 1.75 fish/m³ did not affect pH values ($P > 0.05$). This result is in agreement with Shaker *et al.* (2002). However, the increase in fish density from 1.75 to 14 fish/m³ has significantly affected the pH values indicating that they decreased significantly with the increase in stocking density the pH values increased by several folds in the semi-intensive system compared to that in the extensive one. Dissolved oxygen (DO mg/l) concentration ranged from 4.3 to 6.9 mg/l in all treatments. DO concentrations in extensive ponds did not vary significantly, while the difference was significant between semi - intensive ponds and others. The use of paddle wheel in semi-intensive ponds enhanced significantly ($p < 0.05$) the DO concentrations in ponds furnished by those aerators compared to the rest of the ponds. Results of Table (2) revealed that differences among the applied treatments in averages of water temperature were insignificant and ranged between 27.1 and 27.5°C. The pH, temperature and dissolved oxygen

were the most influencing parameters in fish ponds, where their values in all ponds although fluctuated from time, to time they stayed within the acceptable and favorable levels required for growth, survival and well being of the tested fish species.

The SD readings were higher in the semi-intensive ponds indicating the low abundance of plankton in these ponds compared to the extensive cultured ponds. These results are in agreement with those obtained by Shaker *et al.* (2002).

The average levels of ammonia-nitrogen were 2.5; 2.7; 1.3; 1.4; 1.5 and 1.6 mg/l for semi-intensive treatments (1,2) and extensive treatments (3-6), respectively. The concentrations of the un-ionized ammonia (toxic form) $\text{NH}_3\text{-N}$ in the present study was lower than those recorded in fertilized fish ponds at Lake Burullus (Mosua, 2004). The increase of $\text{NH}_3\text{-N}$ in the semi-intensive ponds compared to other treatments could be explained by the decomposition of organic matter and via the direct excretion of ammonia by the large biomass of fish. The NO_2 and NO_3 concentrations in water followed the same trend of ammonia-nitrogen. The concentrations of NO_2 and NO_3 were also higher in semi-intensive treatments. These results may be due to the consumption of nitrate (which is an essential nutrient) by phytoplankton communities in treatments 3-6. Also, the increase of nitrate in semi-intensive treatments may be related to the decrease of phytoplankton standing corps. It is of particular interest to notice a negative correlation between nitrate content and total phytoplankton which may be attributed to high consumption rate of $\text{NO}_3\text{-N}$ by the dense vegetation. These results are in harmony with those obtained by Shaker *et al.* (2002); Islam (2002) and Mousa (2004). The average concentrations of total alkalinity (T. alk.) and total hardness (T.H) were suitable for fish growth, survival and well being. These results are in agreement with those obtained by Mousa (2004). The average concentrations of total phosphorus (T.P) and orthophosphate (O.P) were lesser in semi-intensive than in the other treatments. These results may be due to the water exchange in semi-intensive pond that led to the decrease of organic matter in these ponds. The average concentration of chlorophyll "a" increased with decreasing fish density per m^3 .

Iron, manganese, zinc, copper, lead, cadmium and mercury were detected monthly during the experimental period in all ponds (Table 3). While the average concentration of mercury did not vary significantly among treatments, these of Fe, Mn, Zn, Cu and Pb varied significantly between semi-intensive ponds and other ponds. It is worthy to mention

that the mean concentrations of heavy metals stayed within the safety range established for aquaculture by FAO (2001), WHO (1989) and Mancy (1993).

Heavy metals concentrations in fish organs are presented in Table (4). The highest levels were found in liver, followed by gills and then muscles in semi-intensive and extensive ponds. These metal concentrations were accumulated from the blood system of fish. In this connection, Shereif and Moaty (1995) found that heavy metals were significantly higher in fish viscera, including liver tissue than in the edible muscles. On the other hand, the highest accumulation of heavy metals in fish species was found in mullet followed by common carp then Nile tilapia and silver carp (Table 4). Generally, it is found that the accumulation level of heavy metals of the present fish species was within the permissible legal limits for different countries (USEPA, 1986 ; WHO, 1989). For example, in the United Kingdom the legal limits for Zn and Pb in fishery products are 50 and 2-10 µg/g, respectively; in Australia it is 40 µg/g for Zn and 1.5-5.5 µg/g for Pb (Nauen, 1983), while the WHO guidelines sets the limits as 1000 µg/g for Zn and 2.0 µg/g for Pb (WHO, 1989). Mosua (2004) recorded concentration levels of iron, zinc, cadmium and lead as 12.2, 30.0, 0.5 and 5.2 µg/g respectively in Nile tilapia in Lake Burullus. However, Noble (1975), Turner *et al.* (1986) and Edwards (1992) stated that there may not be a substantial public health problem produced by the bioaccumulation of toxic elements because almost all organisms contained levels of contaminants, which are below Egyptian National Legal Limits.

As presented in Table (5), the average numbers of phytoplankton and zooplankton decreased with the increase of fish stocking density. The average number of phytoplankton and zooplankton in semi-intensive were 2280 and 99; and 6773 and 380.5 org. /l in extensive ponds, respectively. These results clearly demonstrate that the total amount of phytoplankton and zooplankton decreased in semi-intensive ponds compared to extensive ponds. These results are in agreement with those obtained by Shaker (2002). Semi-intensive farming is usually based solely on artificial feed, while extensive system depends on natural food supplemented by artificial feed which in turn decreases the production cost per kg of fish in the extensive system. Generally, there were positive correlation among total phosphors (T.P); ortho phosphate (O.P); chlorophyll "a" and plankton community and inverse correlation with SD reading in fish

ponds. Chlorophyll "a" contents in water ponds were related to the occurrence of phytoplankton in ponds water.

Fish production and growth performance parameters are illustrated in Tables (6 & 7). The average final weights of Nile tilapia were 216 and 194.25 g for semi-intensive and extensive ponds. Respectively. It is clear that the average weights of Nile tilapia and mullets were higher in semi-intensive system than in extensive treatments. This may be due to the use of artificial feed as the only nutritional source in the semi-intensive system and both species were able to utilize it efficiently. In the natural food medium that supplemented by artificial feed in extensive system, silver carp, common carp and catfish were able to utilize natural communities better and so attaining higher growth rates in extensive treatments than in semi-intensive ones. The average of total fish production for all treatments were 10.374; 13.911; 1.743.5; 2.042.5; 2.142.5 and 2.286.5 kg/feddans for 10; 14; 1.0; 1.26; 1.5 and 1.7 fish/m³ treatments, respectively. These results clearly demonstrate the positive correlation between fish production and stocking density. These findings agree with those obtained by Shaker *et al.* (2002), Sumagaysay and Lourdes (2003) and Yang Yi *et al.* (2003) who found that phytoplankton based food chain was relatively unimportant in pond culture that relies on artificial feed to promote fish growth.

From the data presented in Table (7), it is clear that the daily gain of catfish was higher than other fish species as silver carp, common carp, mullet and Nile tilapia. Also, the daily gain of different fish species was higher in extensive ponds than semi-intensive ponds. These results indicated that the daily gain for each species was negatively correlated with stocking density. Moreover, the same trend was observed in the specific growth rate (SGR). These results are in agreement with those obtained by Karplus *et al.* (1996) and Yang Yi *et al.* (2003). The condition factor did not vary among all species in all treatments ($P < 0.05$).

The economical analysis in the present study is presented in Table (9) and the comparison between semi-intensive and extensive treatments is presented in Table (8). The total values of total return per Feddan were 62.491; 82.809; 9.670; 11.832.5; 12.822 and 14.184.5 LE for T1; T2; T3; T4; T5 and T6. respectively. These results indicated that the highest total return was obtained in semi-intensive treatments. As well as in treatment 2, these indicating a positive correlation between total return and stocking density. Also, the same trend was observed in total costs, where they

increased with increasing stocking density and consequently feed costs. The finding of Shaker *et al.* (2003) and Yang Yi *et al.* (2003) are in agreement with obtained results. Moreover, the same trend was observed by the average cost/kg fish, while the opposite trend was observed in the rate of return to capital as percentage.

REFERENCES

- American Public Health Association (APHA), (2000). Standard Methods for the Examination of Water and Wastewater. The 16th edition, Washington, D. C.
- Edwards, P. (1992). Reuse of human wastes in aquaculture. A technical review. UNEP-World Bank & Sanitation program. The World Bank, Washington DC, USA.
- Food and Agriculture Organization of the United Nations (FAO) (1999). The state of the world fisheries and aquaculture 1998. FAO. Rome, Italy.
- Food and Agriculture Organization of the United Nations (FAO) (2001). Summary tables of fishery statistics; world aquaculture production by principal species. <http://www.fao.org/fi/statist/summtab/fault.asp>
- Hepher, B. (1988). Nutrition of pond fishes. Cambridge University press. Cambridge.
- Islam, S. M.. (2002). Evaluation of supplementary feeds for semi-intensive pond culture of mahseer, *Tor putitora* (Hamilton). *Aqua.*, 212: 263-276.
- Karplus, I.; Milstein, A.; Cohen, S. and Harpaz, S. (1996). The effect of stocking different ratios of common carp, *Cyprinus carpio* L., and tilapias in polyculture ponds on production characteristics and profitability. *Aqua. Res.*, 27: 447-453.
- Lin, H.R. (1982). Polyculture system of freshwater fish in China. *Canadian Journal of Fisheries and Aquatic Sci.*, 39: 143-150.

- Milstein, A. (1992). Ecological aspects of fish species interactions in polyculture ponds. *Hydrobio.*, 231: 177-186.
- Mancy, K.H. (1993). Wastewater treatment at Suez experimental station progress report. National Institute of Oceanography and Fisheries.
- Moussa, S. M. S. (2004). Impact of inorganic pollutants on aquatic environment and fish performance in lake Borollus. Ph. D. Thesis, Inst. of Environment Stud. and Res., Biol. & Phys. Dep., Ain hams Univ., Egypt.
- Noble, R. (1975). Growing fish in sewage. *New Sci.* 67: 259-261.
- Nauen, C. E. (1983). Compilation of legal limits for hazardous substances in fish and fishery products. FAO Fisheries Circular No. 764., Rome, Italy, pp.
- Shaker, I. M. A.; El-Nagdy, Z. A and Ibrahim, N. A. (2003). Effect of organic manure and artificial feeding on water quality and growth performance on Mullet in earthen ponds in Sahl El-teena, Sinai, Egypt. The 1st Scientific conference of the Egyptian Aquaculture Society, Fac. of Environ. Agric. Sci., El-Arish, North Sinai, Egypt, 13-15 Dec. pp.77-96.
- Shaker, I. M. A.; Ibrahim, N. A.; Dawa, M. A. A. and Zakar, A. H. (2002). Effect of stocking density on water quality and mullet growth in earthen ponds at Sahl El-Teena - Senai - Egypt. The 6th Vet. Med. Zagazig. Conference, 7-9 Sept. 2002, Hurghada, Egypt, pp.
- Shereif, M.M. and Moaty, M.A.(1995). Concentration levels and accumulation metals in Nile tilapia grown in wastewater fed fish ponds. In the second Middle East Conference for Waste Management Cairo Publications. Egypt. pp. 284-298.
- Statistical Analysis System (SAS) (2000). SAS program Ver 6.12, SAS Institute Incorporation, Cary, NC 27513, USA.
- Steel, R.G.D and Torrie, J.H (1960). Principles and procedures of statistics. McGraw-Hill Co., New York.

- Sumagaysay, C. N. S. and Lourdes, S. D. M. (2003). Water quality and holding capacity of intensive and semi-intensive milk fish (*Chanos chanos*) ponds. *Aqua.*, 219: 413-429.
- Turner, J.W.D.; Sibbald, R.R. and Hemens, J. (1986). Chlorinate secondary domestic effluent as a fertilizer for marine aquaculture. 3- Assessment of bacterial and viral quality and accumulation of heavy metals. *Aqua.*, 53: 157-168.
- U. S. E. P. A. (Environmental Protection Agency (1986). Quality Criteria for Water. 44015-86-001 Washington.
- WHO (1989). Health guidelines for the use of the wastewater in agriculture and aquaculture. WHO Technical Report Series, No. 778., World Health Organization, Geneva.
- Yang Yi; Kwei, C. L. and Diana, J. S. (2003). Hybrid Catfish (*Clarias macrocephalus* x *C. gariepinus*) and Nile tilapia (*Oreochromis niloticus*) culture in an integrated pen-cum pond system: growth performance and nutrient budgets. *Aqua.*, 217: 395-408.

Table(1): Stocking fish number and weight for different fish species in experimental ponds.

Fish Sp. Treat	Nile Tilapia			Mullet			Common Carp			Silver Carp			Catfish			Total		No. of fish per-m ³
	No.	I.W g	T.w kg	No.	I.W g	T.w kg	No.	I.W g	T.w kg	No.	I.W g	T.w kg	No.	I.W g	T.w kg	No.	T.w	
1	39550	40 ± 5.0	1582 ± 15.75	2000	50 ± 3.0	100 ± 5.5	600	50 ± 3.0	30.0 ± 1.2	200	50 ± 5.0	10.0 ± 0.7	200	50 ± 5.0	10.0 ± 0.5	4255 0	1732 ± 31.5	10.1
2	55050	40 ± 5.0	2202 ± 16.65	3000	50 ± 3.0	150 ± 6.5	800	50 ± 5.0	40.0 ± 1.3	200	50 ± 5.0	10.0 ± 0.7	200	50 ± 5.0	10.0 ± 0.5	5905 0	2412 ± 35.5	14.1
3	3525	10 ± 2.0	35.25 ± 3.4	250	20 ± 1.5	5.0 ± 0.4	125	30 ± 3.0	3.75 ± 0.2	200	50 ± 5.0	10.0 ± 0.7	200	50 ± 5.0	10.0 ± 0.5	4200	64.0 ± 3.2	1.0
4	3875	10 ± 2.0	38.75 ± 3.4	750	20 ± 1.5	15.0 ± 0.75	400	30 ± 3.0	12.0 ± 0.5	200	50 ± 5.0	10.0 ± 0.7	200	50 ± 5.0	10.0 ± 0.5	5325	85.75 ± 3.4	1.268
5	4434	10 ± 2.0	44.34 ± 3.55	1250	20 ± 1.5	25.0 ± 1.1	400	30 ± 3.0	12.0 ± 0.5	200	50 ± 5.0	10.0 ± 0.7	200	50 ± 5.0	10.0 ± 0.5	6384	101.3 4 ± 3.44	1.52
6	4575	10 ± 2.0	45.75 ± 3.75	1750	20 ± 1.5	35.0 ± 1.2	800	30 ± 3.0	24.0 ± 0.75	200	50 ± 5.0	10.0 ± 0.7	200	50 ± 5.0	10.0 ± 0.5	7425	124.7 5 ± 5.5	1.77

Table (2): Total annual means of some physico-chemical characteristics of water samples on experimental ponds.

Items Treat.	Density Fish/ m ³	Temp. °C	pH	D. O mg/l	Satura. %	SD Cm	NH ₃ mg/l	NO ₂ mg/l	NO ₃ mg/l	T. N mg/l	E. C mmhos/Cm	Salinity g/l	T. alk mg/l	T. H mg/l	T. P mg/l	O. P mg/l	Chlorophyll "a" µg/l
1	10.1	27.5 ± 1.1 a	8.6 ± 0.9 b	6.7 ± 1.0 a	82.5 ± 5.5 a	25.5 ± 2.5 a	2.5 ± 0.4 a	0.37 ± 0.12 b	0.36 ± 0.1 b	5.5 ± 1.6 a	2.5 ± 0.3b	2.0 ± 0.1 b	290 ± 50 b	410 ± 72 b	0.92 ± 0.06b	0.32 ± 0.06b	24.5 ± 4.6 b
2	14.1	27.4 ± 0.7 a	8.45 ± 1.0 b	6.9 ± 1.0 a	86.1 ± 5.1 a	27.5 ± 3.5 a	2.7 ± 0.6 a	0.46 ± 0.11	0.42 ± 0.12a	5.9 ± 1.8 a	2.5 ± 0.2 b	1.9 ± 0.1 b	295 ± 42 b	435 ± 70 b	0.96 ± 0.14b	0.34 ± 0.07b	27.5 ± 3.1 b
3	1	27.1 ± 0.6 a	9.5 ± 1.1 a	4.3 ± 0.5 b	65.7 ± 3.5 b	10.5 ± 1.0 b	1.3 ± 0.4 b	0.12 ± 0.03 c	0.16 ± 0.05c	3.3 ± 1.1 b	3.3 ± 0.3 a	2.6 ± 0.1 a	375 ± 64 a	485 ± 56 a	1.56 ± 0.24a	0.76 ± 0.11a	114.6 ± 12.6 a
4	1.268	27.2 ± 0.8 a	9.6 ± 0.9 a	4.5 ± 0.5 b	66.4 ± 6.5 b	12.0 ± 1.0 b	1.4 ± 0.5 b	0.14 ± 0.04 c	0.16 ± 0.05c	3.25 ± 0.9 b	3.2 ± 0.3 a	2.6 ± 0.1 a	380 ± 60 a	486 ± 66 a	1.72 ± 0.17a	0.77 ± 0.14a	96.5 ± 14.1 a
5	1.52	27.2 ± 1.0 a	9.55 ± 0.5 a	4.4 ± 0.5 b	64.5 ± 4.5 b	12.5 ± 1.5 b	1.5 ± 0.5 b	0.16 ± 0.04	0.19 ± 0.06c	3.36 ± 1.1 b	0.31 ± 0.3 a	2.6 ± 0.1 a	380 ± 56 a	505 ± 54 a	1.74 ± 0.19a	0.76 ± 0.13a	96.5 ± 10.1 a
6	1.77	27.2 ± 0.6 a	9.56 ± 0.5 a	4.4 ± 0.4 b	65.1 ± 6.1 b	14.0 ± 1.5 b	1.6 ± 0.3 b	0.22 ± 0.05 c	0.25 ± 0.06c	3.4 ± 1.16b	3.0 ± 0.3 a	2.7 ± 0.1 a	385 ± 52 a	495 ± 74 a	1.76 ± 0.21a	0.75 ± 0.19a	99.5 ± 6.9 a

Means in the same column followed by different letters are significantly different (Duncan s Multiple Range Test P<0.05).

Temp. = Temperature °C, D.O =Dissolved Oxygen, Satura. = saturation%, T.alk. = total alkalinity, T.H = total hardness, T.P = total phosphorus, O.P = ortho phosphate

Table (3): Total annual means of some heavy metals in water samples of experimental ponds as ppm.

Items Treat.	Density Fish/ m ³	Fe	Mn	Zn	Cu	Pb	Cd	Hg
1	10.1	0.5b ± 0.04	0.24b ± 0.09	0.22b ± 0.02	0.26ab ± 0.04	0.14b ± 0.02	0.075b ± 0.01	0.094a ± 0.01
2	14.1	0.52b ± 0.03	0.24b ± 0.09	0.22b ± 0.04	0.24b ± 0.06	0.15b ± 0.02	0.074b ± 0.007	0.09a ± 0.01
3	1.0	0.62a ± 0.03	0.32a ± 0.07	0.28a ± 0.03	0.31a ± 0.09	0.21a ± 0.03	0.12a ± 0.02	0.09a ± 0.01
4	1.268	0.62a ± 0.03	0.34a ± 0.06	0.27a ± 0.02	0.3a ± 0.07	0.22a ± 0.03	0.14a ± 0.022	0.106a ± 0.011
5	1.52	0.65a ± 0.06	0.33a ± 0.07	0.28a ± 0.03	0.26ab ± 0.09	0.24a ± 0.04	0.11a ± 0.04	0.118a ± 0.021
6	1.77	0.64a ± 0.05	0.33a ± 0.09	0.26a ± 0.05	0.27ab ± 0.09	0.23a ± 0.04	0.13a ± 0.01	0.107a ± 0.034

Means in the same column followed by different letters are significantly different (Duncan s Multiple Range Test P<0.05).

Table (4): Average concentrations of some heavy metals ($\mu\text{g/g}$) in liver, gills and muscles tissue for fishes species.

Fish organs		Nile Tilapia				Mullet				Common Carp				Silver Carp			
		Fe	Zn	Cd	Pb	Fe	Zn	Cd	Pb	Fe	Zn	Cd	Pb	Fe	Zn	Cd	Pb
1	Liver	110.24	170.56	6.46	72.82	111.65	172.46	6.78	74.42	110.76	171.08	6.58	72.96	105.46	160.42	5.72	70.06
	Gills	62.8	40.2	5.54	60.2	64.06	40.56	6.02	61.08	63.12	40.44	5.54	60.92	60.72	36.76	5.12	54.98
	Muscle	12.64	34.72	0.56	5.22	14.76	36.5	0.72	6.1	13.2	35.24	0.61	5.56	11.22	30.42	0.46	4.82
2	Liver	115.86	176.12	6.92	75.06	116.66	176.7	6.96	75.01	115.02	172.96	6.96	73.09	109.06	162.92	5.96	70.7
	Gills	66.1	44.16	5.7	62.08	66.7	44.22	6.19	62.0	65.62	42.06	5.88	61.14	61.92	37.22	5.7	55.19
	Muscle	12.7	34.96	0.58	5.26	15.05	36.82	0.86	6.18	13.32	35.61	0.66	5.71	11.28	30.9	0.46	4.86
3	Liver	92.8	155.0	5.72	70.62	93.42	156.12	5.86	71.04	92.96	155.46	5.86	70.9	90.92	142.12	4.56	66.6
	Gills	54.2	35.1	4.9	58.92	54.68	35.26	4.96	59.42	54.46	35.2	4.92	59.04	50.06	32.22	4.5	52.62
	Muscle	9.66	29.46	0.36	4.74	9.76	29.58	0.39	4.92	9.7	29.5	0.38	4.82	9.22	26.44	0.35	4.02
4	Liver	100.04	155.06	5.66	71.02	100.92	155.72	5.74	71.46	99.92	156.52	5.84	71.76	95.72	144.42	4.72	67.44
	Gills	56.72	36.62	4.9	59.66	56.96	37.32	5.24	60.24	56.42	36.78	5.16	59.82	50.92	33.2	4.62	53.0
	Muscle	10.42	32.72	0.4	4.96	10.68	33.34	0.48	5.22	10.5	33.52	0.45	5.12	9.62	26.82	0.36	4.3
5	Liver	101.42	157.75	5.78	70.7	101.94	159.02	6.02	71.04	101.24	158.42	5.96	70.96	96.45	146.9	5.12	68.4
	Gills	57.06	38.8	4.96	59.06	57.64	38.98	5.2	60.16	57.46	38.92	5.1	59.56	52.46	34.4	4.72	53.26
	Muscle	11.06	33.52	0.42	5.02	11.18	34.42	0.52	5.14	10.6	33.92	0.48	5.1	9.9	27.26	0.38	4.42
6	Liver	105.2	160.22	6.01	71.18	106.14	160.92	6.42	71.48	105.7	160.44	6.22	71.22	100.72	150.96	5.42	69.12
	Gills	59.2	39.24	5.12	59.92	59.76	39.46	5.32	60.12	59.3	39.46	5.24	60.02	54.62	35.02	4.88	54.02
	Muscle	11.42	34.04	0.5	5.12	11.66	34.42	0.54	5.36	11.5	34.12	0.52	5.22	10.02	28.24	0.42	4.56

Table (5): Average counts and identification of phytoplankton and zooplankton in the experimental ponds.

Items Treat.	Phytoplankton (org $\times 10/L$)					Zooplankton (org/L)				
	Blue-Green	Green	Bacillaroph.	Cyanoph.	Total	Copep.	Cladocera	Rotifera	Ostracoda	Total
1	1056	876	538	112	2582	51	31	19	11	112
2	872	672	356	78	1978	42	24	11	9	86
3	4566	3250	1110	818	9744	197	190	94	51	532
4	2976	2116	918	716	6726	136	117	88	45	386
5	2578	1750	798	696	5822	124	92	69	41	326
6	2096	1450	678	576	4800	105	81	57	35	278

Bacillaroph.=Bacilliarophyta , Cyanoph = Cyanophyta

Table (6): Fish production and net production of experimental ponds.

Fish sp. Treat.	Nile Tilapia		Mullet		Common Carp		Silver Carp		Catfish		Total production kg	Net production kg
	A.Wt g	T.Wt kg	A.Wt g	T.Wt kg	A.Wt g	T.Wt kg	A.Wt g	T.Wt kg	A.Wt g	T.Wt kg		
1	225 ± 26a	8899 ± 356b	250 ± 26a	475 ± 22b	900 ± 101b	540 ± 26b	550 ± 30c	110 ± 10c	1750 ± 156a	350.0 ± 42a	10374 ± 356b	8642 ± 304b
2	220 ± 26a	12111 ± 456a	250 ± 36a	650 ± 26a	875 ± 110b	700 ± 36a	500 ± 30c	100 ± 10c	1750 ± 166a	350.0 ± 32a	139111 ± 356a	11499 ± 324a
3	240 ± 29a	846 ± 99c	350 ± 22a	85.0 ± 16e	1700 ± 130a	212.5 ± 18c	1400 ± 76a	280 ± 14a	1600 ± 165a	320.0 ± 22a	1743.5 ± 372c	1679.5 ± 316c
4	240 ± 36a	930 ± 91c	270 ± 38a	192.5 ± 19d	1050 ± 109b	420 ± 20b	1150 ± 75b	230 ± 12b	1350 ± 172ab	270.0 ± 20b	2042.5 ± 136c	1956.75 ± 136c
5	230 ± 30a	1021 ± 79c	250 ± 29a	259.5 ± 28d	980 ± 92b	392 ± 14b	1050 ± 64b	210 ± 12b	1300 ± 148ab	260.0 ± 30b	2142.5 ± 118c	2041.16 ± 112c
6	230 ± 20a	1052 ± 109c	250 ± 16a	380.5 ± 29c	590 ± 75c	472 ± 16b	810 ± 44bc	162 ± 16c	1100 ± 120b	220.0 ± 18b	2286.5 ± 98c	2161.75 ± 88c

Means in the same column followed by different letters are significantly different ($P < 0.05$).

Table (7) : Growth performance of different fish species under different aquaculture systems on the experimental ponds.

Items Treat.	Daily Gain					Specific Growth Rate					Condition factor				
	N.Tilapia	Mullet	C.Carp	S.Carp	Catfish	N.Tilapia	Mullet	C.Carp	S.Carp	Catfish	N.Tilapia	Mullet	C.Carp	S.Carp	Catfish
1	1.23 ± 0.14a	1.47± 0.11b	5.7 ± 0.24b	3.33± 0.36c	11.47± 1.92a	1.15 ± 0.1b	1.41± 0.1c	1.93± 0.12b	1.59 ± 0.12b	2.37 ± 0.2a	1.39 ± 0.2a	1.2 ± 0.11a	2.5 ± 0.15a	2.51 ± 0.14a	2.57 ± 0.22a
2	1.2 ± 0.1a	1.47± 0.11b	5.5 ± 0.46b	3.0 ± 0.26b	11.47± 1.56a	1.14 ± 0.1b	1.41± 0.1c	1.91± 0.16b	1.54 ± 0.12b	2.37 ± 0.22a	1.56 ± 0.17a	1.22 ± 0.11a	2.67 ± 0.15a	2.54 ± 0.12a	2.67 ± 0.24a
3	1.53 ± 0.13a	2.13± 0.12a	11.13 ± 1.1a	9.0 ± 1.16a	10.33± 1.11a	2.12 ± 0.12a	1.64± 0.1a	2.69± 0.17a	2.22 ± 0.12a	2.31 ± 0.17a	1.45 ± 0.14a	1.3 ± 0.12a	2.64 ± 0.12a	2.76 ± 0.11a	2.69 ± 0.21a
4	1.53 ± 0.15a	1.6 ± 0.12b	6.8 ± 0.46b	7.47± 0.46b	8.67 ± 1.1b	2.12 ± 0.12a	1.47± 0.1b	2.4 ± 0.17a	2.09 ± 0.4a	2.19 ± 0.21a	1.59 ± 0.12a	1.2 ± 0.1a	2.57 ± 0.14a	2.52 ± 0.32a	2.46 ± 0.17a
5	1.47 ± 0.14a	1.47± 0.12b	6.2 ± 0.76b	8.33± 0.96b	8.33 ± 0.96b	2.09 ± 0.14a	1.41± 0.1b	2.33± 0.17a	2.03 ± 0.17a	2.17 ± 0.17a	1.56 ± 0.13a	1.23 ± 0.1a	2.62 ± 0.12a	2.54 ± 0.24a	2.57 ± 0.24a
6	1.47 ± 0.14a	1.47± 0.12b	5.07± 0.39b	7.0 ± 0.79b	7.0 ± 0.79b	2.09 ± 0.15a	1.41± 0.1b	1.99± 0.17b	1.99 ± 0.17b	2.06 ± 0.15a	1.56 ± 0.12a	1.17 ± 0.1a	3.12 ± 0.11a	2.59 ± 0.26a	2.57 ± 0.14a

Means in the same column followed by different letters are significantly different (P<0.05).

Table (8): Effect of production system on extensive and semi intensive fish culture.

treatments Items	Extensive	Semi-intensive
Stocking denisity/m ³	1.39	12.1
Fish production	2053.75	12142.5
Net production	1959.79	10070.5
Total phytoplankton org/l	6773	2280
Total zooplankton	380.5	99
Daily gain	25.68	22.92
SGR	10.17	8.41
Condition factors	10.7	10.4
Total costs	6098.35	59579
Net return	6028.9	13071
Average cost/kg	5.88	4.865
Return kg above T.V.C. LE	2.94	1.12
Rate of return to capital %	200.73	123.3

Table(9): Economic efficiency for different for aquaculture system in earthen ponds.

Trea Items	Treatment1		Treatment2		Treatment3		Treatment4		Treatment5		Treatment6	
	Quant	LE	Quant	LE	Quant	LE	Quant	LE	Quant	LE	Quant	LE
T.1 st class	4741	33187	6259	43813	476	3332	546	3822	559	3913	570	3990
T. 2 nd class	3072	15360	4079	20395	292	1460	306	1530	336	680	350	1750
T. 3 rd class	966	3864	1346	5384	66	264	66	264	112	448	116	464
T. 4 th class	120	240	427	854	12	24	12	24	14	28	16	32
M. 1 st class	405	4860	521	6252	85	1020	170	2040	236.5	2838	296	3552
M. 2 nd class	70	630	129	1161			25.5	202.5	23.0	207	84.5	760.5
C.C	540	2160	700	2800	212.5	850	420	1680	392	1568	472	1888
S.C	110	440	100	400	280	1120	230	920	210	840	162	648
CF	350	1750	350	1750	320	1600	270	1350	260	1300	220	1100
Total prod.	10374	62491	13911	82809	1743.5	9670	2042.5	11832.5	2142.5	12822	2286.5	14184.5
T. 40g	39.55	9887.5	55.05	13762.5	0	0	0	0	0	0	0	0
T. 10g	0	0	0	0	3525	352.5	3875	387.5	4434	443.4	4575	457.5
M. 50g	2000	700	3000	1050	0	0	0	0	0	0	0	0
M. 20g	0	0	0	0	250	62.5	750	187.5	1250	312.5	1750	437.5
CC 30g	600	60	800	80	125	12.5	400	40	400	40	800	80.0
SC 40g	200	20	200	20	200	20	200	20	200	20	200	20.0
Cf 100g	200	70	200	70	200	70	200	70	200	70	200	70.0
T. C.F	0	10737.5		14982.5		517.5		705		885.9		1065
Feed	18.15	36300	27.6	55200	1.8	3600	2.4	4800	2.75	5500	3.0	6000
Manure	0	0	0	0	1.2	60	1.2	60	1.2	60	1.2	60
Urea	0	0	0	0	0.154	6.5	0.154	6.5	0.154	6.5	0.154	6.5
Superphos	0	0	0	0	0.615	13.5	0.615	13.5	0.615	13.5	0.615	13.5
Aerator	0	519	0	519	0	0	0	0	0	0	0	0
Labor	0	250	0	250	0	50	0	50	0	50	0	50
T.V.C	0	47806.5	0	70951.5	0	4247.5	0	5635	0	6515.9	0	7195
F. Cost	0		0	0	0	0	0	0	0	0	0	0
Dep. P.	0	200	0	200	0	200	0	200	0	200	0	200
T. costs	0	48006.5	0	71151.5	0	4447.5	0	5835	0	6715.9	0	7395
Net R.	0	14484.5	0	11657.5	0	5222.5	0	5997.5	0	6106.1	0	6789.5
Av. C.	0	462	0	511	0	255	0	286	0	313	0	323
Av. Pri.	0	602	0	595	0	555	0	579	0	598	0	620
R.A.T.V.C (LE)	0	140	0	84	0	300	0	293	0	285	0	297
RRC %	0	130.2	0	116.4	0	217.4	0	202.8	0	190.9	0	191.8

F.Prod. – Fish production.

C.C- Common carp

S.C.- Silver Carp

CF- Catfish

T. Prod.- Total production

T- Tilapia

M- Mullet

T.C.F- Total costs of fish

Superphos.- superphosphat

F.Cost- Fixed cost

Dep. P.- Deprecation pond

T. Costs- Total cost

Av. C. – Average Cost/kg

Av. Pri.- Average price

R.A.T.V.C- Return Above T.v.c

- F.Prod. : Fish production .

- C.C : Common Carp .

- S.C : Silver Carp.

- Cf : Catfish.

- T. prod. : Total production.

- V. costs : Variable costs.

- Finger. : Fingerlings.

- T. : Tilapia.

- M. :Mullet.

- F. Cost :Fixed costs.

- Dep. P. : Deprecation pond.

- T. costs: Total cost

- Net R. : Net return.

- Av. C.: Average cost /kg.

- Av. Pri.: Average price.

- R.abv.: Retun above.

- RRC %: Rate of return to capital %