

Biodiversity and Distribution of Zooplankton and Macrobenthic Invertebrates Communities in Lake Nasser, Egypt

Marian, G. Nassif

National Institute of Oceanography and Fisheries, NIOF, Egypt
george.marian@hotmail.com

ARTICLE INFO

Article History:

Received: Sept. 30, 2021

Accepted: Oct.26, 2021

Online: Nov. 19, 2021

Keywords:

Lake Nasser,
Zooplankton,
Macrobenthos,
Invertebrates,
Biodiversity,
Long-term Changes

ABSTRACT

Lake Nasser is assumed to be the largest man-made lake in northern Sudan and southern Egypt. It has paramount importance to Egypt as it supplies water to the country. That's why the lake's water quality must be continuously monitored to cope with the difficulties of water scarcity and a reliable water supply. Thus, in March and July 2019, fifteen sites were chosen to reflect the various habitats of the whole lake. Twenty-one adult zooplankton species were found, belonging to Rotifera, Cladocera, and Copepoda with a total density of 16945 org. m⁻³. Copepoda emerged as the dominant constituent of the community both before and during the flood seasons. The zooplankton Shannon Weaver index value was at its highest in the western sector of Tushka ($H' = 2.46$), while the eastern sector of Aswan station had the lowest value of biodiversity ($H' = 1.97$). Macroinvertebrate fauna consisted of 9 species represented by three phyla; Annelida, Arthropoda, and Mollusca with a percentage of 74.62%, 15.41%, and 9.97%, respectively and the total density was 863 org. m⁻² where the highest densities were found in the southern sectors of the lake. The western station of EL Madiq sector revealed the highest benthic Shannon Weaver index ($H' = 1.55$) which could be ascribed to the preponderance of 4 species of arthropods and exclusively prevalence of nymphs of *Ischneura*. Contrary, the middle stations of Wadi Abyad and Aswan sectors recorded the least diversity Shannon Weaver values ($H' = 0.21$ and 0.27 , respectively). The present study revealed that the number of macroinvertebrate species decreased dramatically to 9 compared to 15, 43, 39, and 24 in 1987, 1993, 1995, and 2013, respectively. It is recommended to force an effective legal framework to mitigate pollution in Lake Nasser.

INTRODUCTION

Lake Nasser was built by the construction of the Aswan High Dam between January 1964 and June 1968 (Abd El-Monsef *et al.* 2015; El Gamal & Zaki 2017; Salih *et al.* 2019). The Lake's surface area is approximately 5000 km² (Farhat & Aly 2018). Its storage capacity is 150–165 km³, and supplies a water flow of 11,000 m³/s. The mean depth of Lake Nasser is 90 m, and the maximum width of the Lake is about 60 km (Abou El Ella & El Samman 2010; El Shemy 2010; Khalifa *et al.* 2015).

Zooplankton has a vital role as bio-indicators of water quality. They are significantly influenced by environmental factors and demonstrate rapid responses to variations in environmental conditions (**Gannon and Stemberger, 1978**). Among these organisms, rotifers, with their high population turnover rates, serve as effective indicators of saprobity. Not only can their community structure be utilized to estimate population levels, but it can also provide insights into the overall trend of environmental conditions over time. Furthermore, rotifers play a crucial role in regulating the transfer of nutrients and energy to higher trophic levels (**El-Shabrawy, 2001**). Recent studies by **Khalifa et al. (2015)**, **Salem (2021)**, and **Hegab et al. (2021)** have investigated zooplankton abundance in Lake Nasser.

Latif (1974) and **Iskaros (1988, 1993)** reported that macrobenthic invertebrates are significant food sources of a variety of fish species in Lake Nasser. Macrobenthic invertebrates are better bio-indicators for comprehending the changing aquatic conditions compared to the microbiological and chemical data, which only provide short-term variations (**Ravera, 1998 & 2000**). However, information on invertebrates of Lake Nasser is very rare. **Iskaros (1988 & 1993)** studied in detail the spatial distribution and temporal variations of the macrobenthic animals of the lake where he identified 40 species represented by Aquatic Insecta, Mollusca, Annelida, and Platyhelminthes. Thirty-nine macrobenthos species were recorded by **Fishar (1995)**. **El-Shabrawy & Abd El-Regal (1999)** mentioned that the main channel harbored little diversity; only nine species of Arthropoda, Annelida, and Mollusca were recorded. **Iskaros and Gindy (2009)** studied the impact of substrate status on the benthic fauna in Aswan reservoir during spring, summer, autumn 2008, and winter 2009. **Mola and Abdel-Gawad (2014)** studied the spatial and temporal variations of macrobenthic fauna in Lake Nasser khors during 2013 and they found that Khor Tushka West recorded the highest population density while Khor Wadi Abyad was the lowest. Furthermore, they found that spring exhibited the highest population density compared to the other seasons. Also in 2013, **Abdel-Gawad and Mola (2014)** recorded 24 macrobenthic invertebrate species in Lake Nasser main stream. In addition, **Abdel-Gawad (2016)** studied the impact of physico-chemical factors on the distribution and diversity of molluscs in Lake Nasser and she recorded ten species of molluscs, seven of them were Gastropod while three were Bivalve. Moreover, **Wahab et al. (2018)** studied the community structure, abundance, and diversity of macrobenthic invertebrates in the four northern khors of Lake Nasser during 2015. They recorded 26 species and mentioned that the western khors were higher in species richness and abundance than that of the eastern khors. **Abdel Gawad and Abdel-Aal (2018)** studied the community structure of both phytoplankton and macrobenthic invertebrates associated with macrophyte *Myriophyllum spicatum* in Dahmeit and Tushka west khors of Lake Nasser during pre-flood, flood and post-flood seasons in 2016.

Obviously, there is not enough data on the mainstream of Lake Nasser's zooplankton and bottom fauna, which are considered the main components of the food web and a source of fish diet. Thus, this investigation was dedicated to monitor the zooplankton and macrobenthic invertebrate groups inhabiting the mainstream of Lake Nasser (abundance, distribution, and diversity in association with various physical and chemical parameters and seasonal variation). Additionally, it aimed to spotlight the impact of the flood on the fauna.

MATERIALS AND METHODS

The study area:

Fifteen stations have been chosen, accounting for various habitats of the whole lake (**Fig. 1**). These stations could be described as five sectors (Aswan High Dam, Wadi Abyad, El-Madiq, Tushka, and Abu Simble). Samples were collected from east, middle, and west of each sector.




Fig.1. Map of Lake Nasser showing the selected sampling sectors

Sampling Procedures:

Samples of water, zooplankton, and macrobenthos have been collected from Lake Nasser in March and July of 2019 (both before and during the flood). The standard procedures for water and wastewater examination were followed while analyzing the samples (**APHA, 2005**).

The multi-probe portable meter has been used to measure water's temperature, EC, pH, and DO in the field. A typical 25cm diameter Secchi disc was used to measure the water transparency. The observations were made in the field, and results were expressed in cm at the distance at which the Secchi disc disappeared.

Zooplankton samples were gathered using a 55-micron mesh-size net at a depth of 5 meters from the subsurface water. These samples were then preserved in a 5% neutral formalin solution for subsequent microscopic examination in the laboratory.

A square Ekman grab sampler with a 225cm² opening area was used to collect the macrobenthos samples. Each station's surface layer of the bottom deposits was collected. The specimens have been washed immediately to remove any sediments or mud via a 500µm mesh net and then stored in plastic containers with 7% formalin. The samples underwent another sieving process via a 500µm mesh net in the lab. The benthic animals were sorted into genera and species using a zoom stereo microscope. Each group was counted and preserved in a bottle with 7% formalin. The collected species have been identified to species level when possible.

Data Treatments and Statistical Analyses:

In order to investigate the relationship between the abiotic and biotic parameters, correlation analysis had been applied to the data via Microsoft Excel. The zooplankton and bottom fauna species diversity index was computed and evaluated to assess the devastating impacts of pollution on the species diversity, and eventually the ecosystem via Primer 5 version 5.2.0. The Bray-Curtiz similarity index was applied to study the similarity of the communities inhabiting the studied stations.

RESULTS AND DISCUSSION

Physical and chemical variables:

Water temperature:

Remarkably, the temperature has a direct effect on the aquatic animals and the chemical and physical characteristics as well (Abdo, 2003). Therefore, global warming has a great negative effect on aquatic organisms' diversity. The big variation in temperature of Lake Nasser water all the year-round is considered a controlling feature associated with a scale of tolerance of species (Mageed & Heikal, 2006). During the current study, there was no notable variation between the stations with an annual average of 25°C (Table 1).

Water transparency:

The light penetration decreases with increasing turbidity. This, in turn, affects many essential processes as photosynthesis. In the current investigation, the northern part of the lake shows the highest transparency, especially in the Aswan sector with a yearly average of 318 cm. The lake's transparency gradually decreased downward to reach its lowest value in the Tushka sector with an annual transparency of 215 cm. No significant effect of water transparency was detected on the distribution of the benthic fauna, as shown in the correlation study (Table 1).

The electrical conductivity (EC):

During the investigation period, the electrical conductivity values represented their highest in the eastern part of the Aswan sector in March (before the flood), with an average of 291 mS/cm. It was noticeable that EC values gradually decreased from the north to the south of the Lake with an average of 259 mS/cm. Mageed and Heikal (2006) recorded similar outcomes. There was no significant correlation between EC values and the macrobenthic invertebrates' biodiversity or distribution.

Hydrogen ion concentration (pH):

According to (WHO, 2003), pH is one of the most crucial functional water quality metrics. During the current investigation, there was no significant difference between stations along the lake. Furthermore, pH values were on alkaline side with an average of 8.6.

Dissolved Oxygen (DO):

The peak value of dissolved oxygen in Lake Nasser was in the Tushka sector (annual DO average of 7 mg/l). Furthermore, the western area of the Tushka sector showed its maximum DO

concentration in March (9.6 mg/l). Conversely, El-madiq and Abu Simble revealed the lowest DO concentration in summer (5.3 mg/l). However, the Wadi Abyad sector displayed the minimum yearly DO (average of 6.4 mg/l). Generally, Lake Nasser showed a good status in DO, as reported by the Egyptian Decree No. 92/2013.

Table 1. Spatio-temporal variation of physico-chemical parameters of Lake Nasser during March and July 2019

	M1EM1MM1W	Aswan	M2EM2MM2W	WadiAbyad	M3EM3MM3W	El-Madiq	M4EM4MM4W	Tushka	M5EM5MM5W	Abu Simble
Temp (°C)										
March	18.9 17.8 17.5	18	19.6 19.9 20.3	20	20.2 20.7 20.1	20	20.9 20.1 20.6	21	21.1 21.3 21.4	21
July	28.2 26.7 26.7	27	30 29.7 30.3	30	31.1 31.1 31.1	31	31.4 31.2 30.9	31	30.6 30.6 31.3	31
Average	24 22 22	23	25 25 25	25	26 26 26	26	26 26 26	26	26 26 26	26
Trans (cm)										
March	400 300 310	337	250 400 390	347	250 240 280	257	190 230 220	213	180 180 185	182
July	250 300 350	300	180 310 220	237	270 200 330	267	220 250 180	217	280 280 305	288
Average	325 300 330	318	215 355 305	292	260 220 305	262	205 240 200	215	230 230 245	235
EC mS/cm										
March	291 286 282	286	285 283 283	284	276 278 285	280	264 261 261	262	254 257 257	256
July	246 250 248	248	247 246 247	247	244 245 250	246	237 238 234	236	240 243 237	240
Average	269 268 265	267	266 265 265	265	260 262 268	263	251 250 248	249	247 250 247	248
pH										
March	8.45 8.39 8.37	8.40	8.61 8.54 8.49	8.55	8.77 8.79 8.77	8.8	8.72 8.79 8.7	8.7	8.85 8.84 8.71	8.8
July	8.49 8.6 8.58	8.56	8.5 8.51 8.52	8.51	8.71 8.66 8.44	8.6	8.55 8.57 8.74	8.6	8.43 8.47 8.56	8.5
Average	8.47 8.50 8.475	8.48	8.568.5258.505	8.53	8.748.7258.605	8.7	8.64 8.68 8.72	8.7	8.64 8.66 8.635	8.6
DO (mg/l)										
March	7.27 8.83 6.25	7.45	6.88 6.95 8.08	7.3	8.62 8 8.67	8.4	7.39 8.9 9.6	8.6	9.28 6.98 8.7	8.3
July	5.85 5.48 6.34	5.89	6.05 5.15 5.38	5.5	5.74 4.9 5.13	5.3	5.2 5.09 5.8	5.4	5.52 5.5 5.02	5.3
Average	6.56 7.16 6.295	6.67	6.47 6.05 6.73	6.42	7.18 6.45 6.9	6.84	6.30 7.00 7.7	7.0	7.4 6.24 6.86	6.8

Zooplankton community structure and density:

A total of twenty-one adult zooplankton species, along with three immature copepod stages, were identified in the main stream of Lake Nasser, belonging to three distinct groups: Rotifera, Cladocera, and Copepoda. Copepoda, as illustrated in **Fig. 2**, was the dominant group, representing 45.45% of the community prior to the flood season and 49.10% during the flood season. Extensive research conducted by **Khalifa et al. (2015)**, **Salem (2021)**, and **Hegab et al. (2021)** confirmed the prevalence of copepods in Lake Nasser.

The average total zooplankton density was estimated at 16945 organisms per cubic meter, with Tushka and Abu Simble exhibiting the highest population density, averaging 6593 and 5563 organisms per cubic meter, respectively. Seasonal variation analysis, as depicted in **Fig. 3**, exhibited higher population density in spring compared to summer, potentially attributed to elevated dissolved oxygen content during the spring season. Furthermore, the high water flow rate during the summer (flood season) had a detrimental impact on the weaker zooplankton.

The Copepoda group primarily comprised *Acartia latisetosa*, *Acanthocyclops* sp., *Thermocyclops neglectus*, and nauplius larvae. Notably, nauplius larvae accounted for approximately 54.88% of total copepods, corroborating findings reported by Amer (2007) and Nassif (2012). It is noteworthy that the eastern section of the Abu Simble station demonstrated the highest copepod density, averaging 21386 org. m⁻³. In contrast, the eastern sector of the EL-Madiq station exhibited the lowest density, averaging 2981 org. m⁻³ during July 2019. In terms of seasonal variation, spring showcased the highest copepod population density, while summer exhibited the least population density, with average densities of 8033 and 7963 org. m⁻³, respectively.

Cladocera constituted approximately 18.10% of the total zooplankton population, while this percentage doubled to 38.62% during summer (Fig. 2). Seven species of Cladocera were identified, with *Bosmina longirostris* being the most dominant, representing 35.89% of the total Cladocera density. This finding is in harmony with the previous studies conducted by Amer (2007), Yousry (2009), and Nassif (2012). Conversely, *Alona intermedia* was found to be the least abundant species in the main channel of Lake Nasser during the study period. The highest cladoceran density was observed in the western sector of the Abu Simble station, while the lowest density was recorded in the western sector of Tushka. Overall, summer exhibited higher population density compared to spring.

The prevalence of Rotifera was found to be 36.45% of the total community in spring, which decreased to one-third of the population during summer (Fig. 2). The eastern sector of Tushka displayed the highest rotifer population density, whereas the western sector of Wadi Abyad had the lowest density of Rotifera. Notably, spring exhibited higher population density compared to summer. Moreover, certain species such as *Keratella quadrata*, *Keratella tropica*, and *Lecane luna* were exclusively recorded during the summer season, possibly attributed to their stenothermal characteristics. Conversely, species including *Brachionus angularis*, *Monostyla lunaris*, *Polyarthra vulgaris*, and *Proalids* sp. were solely recorded during spring. The findings of Hegab et al. (2021) were consistent with the current study, offering further support for the observed seasonal variations in the zooplankton community of Lake Nasser.

Zooplankton diversity index:

The analysis revealed that the highest index values were observed in the western sector of Tushka, followed by the western sector of Abu Simble and the western sector of EL-Madiq station, with corresponding entropy values (H') of 2.46, 2.38, and 2.32, respectively (Table 2). The western sector of Abu Simble also had the highest number of species ($S=18$). This may reflect the high ecosystem quality. Conversely, the middle sector of El-Madiq station exhibited the lowest biodiversity value with an entropy value of 1.65. Notably, the summer season exhibited a higher number of species compared to those found in spring, a result that is in line with the findings reported by Hegab et al. (2021).

The similarity dendrogram revealed the presence of two main clusters. The first cluster was found to be comprised of the middle sector of El-Madiq station, highlighting the distinct structure of the zooplankton community in this specific area. Meanwhile, the second cluster,

comprising two additional clusters, demonstrated the diverse nature of the community in Lake Nasser (**Fig. 4**). The community structure of zooplankton in the eastern sector of Tushka and Abu Simbel exhibited noteworthy similarities. Similarly, the communities in the western sector of these stations displayed a high degree of resemblance.




Fig.2. Zooplankton community composition in Lake Nasser, a, in March 2019; b, in July 2019.




Fig.3. Seasonal variation of zooplankton community in Lake Nasser in March 2019 and July 2019.




Fig.4. Dendrogram illustrating the zooplankton similarity of the 15 studied stations in Lake Nasser.

Table 2. Zooplankton diversity indices of the different stations in the current study.

Station	S	N	d (Richness)	J' (Evenness)	H'(log) (Shannon)	1-lambda Simpson
St.1 E	10	18496	0.76	0.79	1.97	0.81
St.1 M	9	9678	0.80	0.81	2.00	0.82
St.1 W	16	9873	1.16	0.77	2.19	0.85
St.2 E	13	12509	1.00	0.78	2.11	0.84
St.2 M	12	8724	0.95	0.84	2.21	0.87
St.2 W	13	14827	0.99	0.82	2.21	0.85
St.3 E	14	21630	1.03	0.81	2.26	0.87
St.3 M	12	17041	0.91	0.62	1.65	0.70
St.3 W	13	9783	1.09	0.84	2.32	0.87
St. 4 E	15	29000	1.08	0.74	2.10	0.84
St.4 M	12	15781	0.91	0.75	1.98	0.81
St.4 W	15	21401	1.17	0.85	2.46	0.90
St.5 E	16	25215	1.09	0.73	2.08	0.79
St.5 M	10	15877	0.77	0.76	1.88	0.79
St.5 W	18	24343	1.29	0.79	2.38	0.87

Macrobenthic invertebrates:

Community structure and density:

Nine macrobenthic invertebrate species were found during the period of study. The benthic invertebrates were related to three different groups; Annelida, Arthropoda, as well as Mollusca, with percentages of 74.62%, 15.41%, and 9.97%, respectively (Fig. 5). A similar result was recorded by Mola and Abdel Gawad (2014). In the current investigation, the macrobenthos total density was 863 org. m⁻². It was noticeable that the community represented a higher density in the eastern sector of the lake during spring (pre-flood), while the density was higher in the middle sector of Lake Nasser in the summer season (flood season).

The highest population densities were found in the southern part of the lake particularly in Tushka and Abu Simble sector with an average of 1340 and 1065 org. m⁻², respectively (Fig. 6). This result was in coincidence with Abdel Gawad and Mola (2014) which may be ascribed to the bottom characteristics and the food availability as it has been seen as the most crucial features that determine the spatial distribution of the benthic community (Iskaros and El-Dardir, 2010), although, Nkwoji *et al.* (2010) pointed out that low macrobenthos community abundance and diversity was significantly impacted by stress caused by pollutants. In this context, the El-madiq sector revealed the poorest density of population, with a yearly average of 445 org. m⁻²; a finding that matches with that of Abdel-Gawad and Mola (2014) who recorded 127 org. m⁻² in the El-madiq sector during their survey in 2013.

Regarding the seasonal variation, summer (flood season) exhibited higher population density than spring (pre-flood) which was due to the dominance of *Limnodrillus* spp. and *Chironomus* larvae (Fig. 7). This result is in coincidence with Iskaros and Gindy (2009) who referred this result to the strong relationship between the benthic fauna standing crop and the amount of organic carbon and calcium carbonate in the sediment. Furthermore, Latif *et al.* (1979) mentioned that the macrobenthos standing crop increases with the rise in water level (flood season).

Annelida was the most predominant macrobenthic invertebrates during this investigation period with a yearly average of 644 org. m⁻² (**Table 3**). It represented 74.62% of the total macrobenthic community. It was represented only by one species *Limnodrilus* sp.. According to **Iskaros and El-Dardir (2010)**, the oligochaetes dominance in Lake Nasser was due to their adaptability to a variety of habitats and their tolerance to anoxic or low oxygen content conditions. Furthermore, the dominance of *Limnodrilus* Spp. was in coincidence with **Iskaros (1993)**, **Fishar (1995)**, **Iskaros and Dardir (2010)**, and **Mola and Abdel Gawad, (2014)**.

For the distribution of *Limnodrilus* sp., the middle station of Abu simble sector showed the maximum abundance with an yearly average of 1500 org. m⁻². On the other hand, the middle area of El-Madiq sector revealed the minimum density with a yearly average of 75 org. m⁻². By and large, Abu simble and Tushka sectors revealed the highest *Limnodrilus* sp. population density with an annual average of 1065 and 1030 org. m⁻², respectively. With respect to seasonal variation, summer (flood season) had the highest density of population (902 org. m⁻²) while spring (pre-flood season) exhibited the lowest abundance (386 org. m⁻²).




Fig.5. Community structure of macrobenthic invertebrates in Lake Nasser




Fig.6. Spatial variation of macrobenthic invertebrates in Lake Nasser during March 2019 and July 2019




Fig.7. Seasonal variation of macrobenthos in Lake Nasser (Org. m⁻²) in March 2019 and July 2019.

Table 3. Seasonal variation of macrobenthos in Lake Nasser (Org. m⁻²) during March and July 2019

	March	July	Average
Mollusca			
<i>Gyraulusehrenbergi</i>	56	18	37
<i>Melanoidestubercolata</i>	10	24	17
<i>Physaacuta</i>	2	0	1
<i>Succinia</i>	2	46	24
<i>Pisidiumpirothi</i>	0	14	7
Subtotal	70	102	86
Arthropoda			
<i>Chironomus</i> Larvae	44	154	99
<i>Cricotopus</i> sp.	0	34	17
Nymph of <i>Ischneura</i>	0	2	1
Pupa of Chironomidae	28	4	16
Subtotal	72	194	133
Annelida			
<i>Lymnodrillus</i> spp.	386	902	644
Total	528	1198	863

Arthropods were ranked as the second dominant benthic fauna during the period of study with a yearly average of 133 org. m⁻² (Table 3). It composed 15.41% of the total macrobenthic invertebrates' community. It was represented by *Cricotopus* sp., nymph of *Ischneura*, larvae, and pupae of *Chironomus* sp. as shown in Fig. 8.

Regarding spatial distribution, the maximum arthropod population density was represented in the western side of Aswan sector due to the dominance of *Chironomus* larvae with an average of 555 org. m⁻², while the middle station of Wadi Abyad revealed the least density with a mean of 15 org. m⁻².

In terms of seasonal variance, the average arthropod population density during summer (flood season) showed its maximum of 194 org. m⁻², whereas the average density during spring (the pre-flood season) was 72 org. m⁻² (Table 3).

Chironomid larvae serve as the main feeding source for *Mormyrus kannume*, *M. caschive*, and *Chrysichthys auratus* in Lake Nasser as mentioned by **Latif (1974)**; **Iskaros (1988)**, and **Iskaros (1993)**. During the present investigation, *Chironomus* larvae constituted 74.44% of the arthropod density, while 11.47% of the whole benthic community, with a yearly mean of 99 org./m². *Chironomus* larvae ranked as one of the highest population densities during the whole period of investigation. This agrees with the results recorded in the study of **Fishar (2000)**. These larvae prefer to live in the littoral zones of eutrophic and oligotrophic lakes. The dominance of Chironomids in reservoirs is usually an indicator of pollution as they can consume the organic matter and able to survive under oxygen-deprived conditions (**Rosenberg et al. 1984**). In the current study, the largest chironomid density was found in the western section of the Aswan sector with an annual average of 540 org. m⁻². It was noticeable that *Chironomus* larvae disappeared from the middle stations of Wadi Abyad and Abu Simble sectors. This could be attributed to the fact that the abundance and dominance of *Chironomus* larvae are related to macrophytes in order to hide from predators and consume the epiphytic microorganisms as a feeding source (**Iskaros et al. 2011**).

For seasonal variation, summer (the flood season) had the largest density (154 org. m⁻²). Similar results were found in the study of **Iskaros (1988, 1993)**. On the other hand, spring (the pre-flood season) showed the lowest population density (44 org. m⁻²).




Fig.8. Community structure of Arthropoda in Lake Nasser

Mollusks made the third rank of the macrobenthic community, which was composed of five species (**Fig. 9**). *Gyraulus ehrenbergi* was the most dominant mollusk species during March (pre- flood season), while *Succinia* sp. was the most predominant species during July (flood season). By and large, *Gyraulus ehrenbergi* was the dominant species during this study and represented 43.02% of the total mollusk community. For spatial distribution, a significant spatial distribution was detected where the western sector of Tushka revealed the maximum density with a yearly average of 465 org. m⁻². On the other hand, mollusks disappeared totally from the Abu Simble sector during the period of investigation.




Fig.9. Community structure of Mollusca in Lake Nasser


Macrobenthos diversity index:

Because each species in the ecosystem has a particular purpose and maintains freshwater resources, a balanced ecosystem with rich biodiversity increases the ecosystem's productivity. As a result, in contaminated ecosystems, a decline in species diversity is understood. Three stations represented the highest species diversity in Lake Nasser and, therefore, reflect the best environment for the organisms to live in (**Table 4**). The western station of the EL Madiq sector revealed the highest Shannon Weaver diversity index ($H' = 1.55$). It could be ascribed to the prevailing of four arthropod species and nymphs of *Ischneura*. This could be a reflection of the excellent water quality in this area. This result is followed by the eastern part of the Wadi Abyad sector and the western part of Tushka which revealed 1.23 and 1.09 of the Shannon Weaver index, respectively. On the contrary, the middle stations of the Wadi Abyad and Aswan sectors recorded the least diversity Shannon Weaver index value ($H' = 0.21$ and 0.27 , respectively). It may be attributed to the nature of the sediment, which led to the absence of all mollusk species and the presence of one Annelida species (*Limnodrilus sp.*) and one Insecta species (*Chironomus* larvae). However, the dominance of *Limnodrilus sp.* and *Chironomus* larvae with the absence of all other species has been used to indicate poor water quality, which is confirmed by **Brinkhurst (1974)**, who mentioned that the monogeneric assemblage of tubificid worms, with the dominance of *Limnodrilus hoffmeisteri*, indicates that the water stream is somewhat gross organic pollution and eutrophic.

Because the macrobenthic invertebrate community is thought to be the excellent indicator of the biological and environmental quality of the aquatic ecosystem, to examine the relationships between the population abundance and the stations, the similarity index was applied and the dendrogram is illustrated in **Fig. 10**. Obviously, there were 2 clusters of stations. The first cluster was composed of Abu Simble sector, the middle and eastern stations of Tushka sector, and the middle stations of both Aswan and Wadi Abyad sectors. It could be attributed to the factor of low diversity index in these stations. The second cluster was characterized by higher biodiversity mostly in the northeastern and western stations (Aswan, Wadi Abyad, and El-Madiq).

Table 4. The benthic macroinvertebrates diversity indices of the different stations in the current study.

Station	S	N	d (Richness)	J' (Evenness)	H'(log) (Shannon)	1-lambda Simpson
St.1 E	3	180	0.39	0.81	0.89	0.54
St.1 M	2	975	0.15	0.39	0.27	0.14
St.1 W	5	1260	0.56	0.60	0.97	0.56
St.2 E	7	720	0.91	0.63	1.23	0.56
St.2 M	2	270	0.18	0.31	0.21	0.11
St.2 W	5	510	0.64	0.61	0.98	0.48
St.3 E	5	735	0.61	0.64	1.02	0.52
St.3 M	4	270	0.54	0.71	0.98	0.55
St.3 W	6	330	0.86	0.87	1.55	0.76
St. 4 E	5	1740	0.54	0.41	0.65	0.32
St.4 M	3	810	0.30	0.29	0.32	0.14
St.4 W	7	1470	0.82	0.56	1.09	0.54
St.5 E	2	1455	0.14	0.67	0.46	0.29
St.5 M	2	1665	0.13	0.47	0.32	0.18
St.5 W	3	555	0.32	0.38	0.42	0.20

**Fig.10.** Dendrogram illustrating the macrobenthos similarity of the 15 studied stations in Lake Nasser.

Long-term changes of macrobenthic invertebrates' community structure:

Obviously, macrobenthic invertebrates are influenced greatly by complex inter-relationships between biotic and abiotic factors. In the present investigation, oligochaetes were the most abundant group, where the decaying detritus and macrophytes provide the bottom with nutrients that assure good nourishment.

Historically, the Lake's substrate was occupied with extremely large numbers of bivalves. During the stagnant period, these bivalves perished and disappeared. Since 1973, oligochaetes and mussels returned to the shallow water of the khors because of the presence of favorable oxygen conditions (Entz 1976).

As shown in **Table (5)**, the number of species decreased dramatically to only 9 species compared to 15, 43, 39, and 24 in 1987, 1993, 1995, and 2013, respectively. This could be related to anthropogenic impacts.

Furthermore, the standing crop of macrobenthic fauna in Lake Nasser was 863 org. m⁻² compared to 823 org. m⁻² in 1995, and 529 org. m⁻² in 2013 (**Table 5**). Thus, it is clear that Lake Nasser has possessed the same composition of macrobenthic invertebrates since the 1980s till now, although the biodiversity and species richness have been declined to a great extent. **Nassif (2012)** deduced that little species variety and dominance of pollution-tolerant species, such as *Limnodrilus hoffmeisteri* and *Chironomus* larvae, beside the disappearance of sensitive species demonstrate that the water status of Lake Nasser has suffered greatly.

CONCLUSION

In conclusion, the current study revealed that the western sector of Abu Simble station had a high biodiversity index value and the highest number of species, indicating the high ecosystem health of this sector. However, the macrobenthos biodiversity, as well as species richness in Lake Nasser, showed a drastic decline from one year to another. Therefore, it is recommended to apply an efficient legal framework and improve public awareness to maintain Lake Nasser healthy. That is why this paper is an alarming tool for decision-makers and stakeholders to take the right action to mitigate the pollution resources discharging on the most crucial and historical lake in Egypt.

Table 5. The macrobenthic invertebrates list in Lake Nasser recorded by other authors (+ = found, - = not found).

Taxa and species	Elewa (1987)	Iskaros (1988 & 1993)	Fishar (1995)	Abdel-Gawad and Mola (2014)	Current study
Phylum : Cnidaria					
Class : Hydrozoa					
<i>Hydra vulgaris</i> (Pallas, 1766)	-	-	+	-	-
Phylum : Bryozoa					
Class : Phyleclolaemata					
<i>Fredericella sultana</i> (Blumenbach, 1779)	-	-	+	-	-
Phylum : Arthropoda					
Class : Insecta					
<i>Ablabesmyia</i> sp.	-	+	+	-	-
<i>Caenis</i> sp.	-	-	+	-	-
<i>Chironomus</i> sp.	-	+	+	+	+
<i>Circotopus</i> sp.	-	+	+	-	+
<i>Clinotanpus</i> sp.	+	+	-	-	-
<i>Coelotanpus</i> sp.	-	+	+	-	-
<i>Conchopelopia</i> sp.	-	+	-	-	-
<i>Cryptochironomus</i> sp.	+	+	+	+	-
<i>Dicrotendipes modestus</i>	-	+	-	-	-
<i>Einfeldina</i> sp.	-	+	-	-	-
<i>Enallagma</i> sp.	-	-	+	-	-
<i>Gomphus</i> sp.	-	+	-	-	-
<i>Ischnura</i> sp.	-	-	+	+	+
<i>Libellula</i> sp.	-	+	-	-	-
<i>Microchironomus</i> sp.	-	+	-	-	-
<i>Micronecta plicata</i>	-	-	+	-	-

<i>Microtendipes</i> sp.	+	+	+	-	-
<i>Neurocordula</i> sp.	-	-	+	-	-
<i>Nilodorum</i> sp.	-	+	+	-	-
<i>Pelopia</i> sp.	-	+	-	-	-
<i>Plathemis</i> sp.	-	-	+	-	-
<i>Polypedilum</i> sp.	-	+	+	+	-
<i>Perithemis</i> sp.	-	+	+	+	-
<i>Procladius</i> sp.	+	+	+	-	-
<i>Pseudoagrion niloticus</i>	-	+	-	-	-
<i>Tanpus</i> sp.	-	+	-	-	-
<i>Dytiscide</i> sp.	-	-	+	-	-
<i>Tanytarsus</i> sp.	+	+	+	+	-
<i>Hydrovatus</i> sp.	-	-	+	+	-
Larvae of Trichoptera	+	+	+	+	-
Pupae of Chironomidae	-	-	-	+	+
Nymphs of Ephemeroptera	-	+	-	-	-
Adult of Corixidae	-	+	-	-	-
Class : Crustacea					
<i>Cardina nilotica</i> (P. Roux, 1833)	-	-	+	-	-
<i>Chlamydotheca unispinosa</i> (Baird, 1862)	-	-	+	+	-
<i>Potamonautes niloticus</i> (H. Milne Edwards)	-	-	-	-	-
<i>Stenocypris malcolmsoni</i> (Baired, 1862)	-	-	+	-	-
Phylum : Annelida					
Class : Oligochaeta					
<i>Branchiura sowerbyi</i> (Beddard, 1892)	+	+	+	+	-
<i>Limnodrilus hoffmeisteri</i> (Claparède, 1862)	+	+	+	+	+
<i>Limnodrilus udekemianus</i> (Claparède, 1862)	+	+	+	+	-
<i>Pristina</i> sp.	-	-	+	+	-
Class : Hirudinea					
<i>Helobdella conifera</i> (Moore, 1933)	-	+	+	+	-
Phylum: Mollusca					
Class : Gastropoda					
<i>Bellamyia unicolor</i> (Olivier, 1804)	-	+	-	-	-
<i>Biomphalaria alexandrina</i> (Ehrenberg, 1831)	-	+	-	-	-
<i>Bulinus truncatus</i> (Audouin, 1827)	+	+	+	+	-
<i>Bulinus forskalii</i> (Ehrenberg, 1831)	-	+	-	-	-
<i>Cleopatra bulimoides</i> (Olivier, 1804)	-	+	+	+	-
<i>Gabbiella senaariensis</i> (Kuster, 1852)	-	+	-	-	-
<i>Helisoma duryi</i> (Wetherbg, 1879)	-	+	-	-	-
<i>Lanistes carinatus</i> (Olivier, 1804)	-	+	-	-	-
<i>Lymnaea natalensis</i> (Krauss, 1848)	-	+	-	-	-
<i>Melanoides tuberculata</i> (Müller, 1774)	+	+	+	+	+
<i>Physa acuta</i> (Darparnaud, 1805)	+	+	+	+	+
<i>Pila ovata</i> (Olivier, 1804)	-	+	-	-	-
<i>Segmentorbis angustus</i> (Jickeli, 1874)	-	+	-	-	-
<i>Theodoxus niloticus</i> (Reeve, 1856)	-	+	-	-	-
<i>Valvata nilotica</i> (Jickeli, 1874)	+	+	+	+	-
<i>Gyraulus ehrenbergi</i> (Beck, 1837)	-	-	+	+	+

Class : Bivalvia					
<i>Corbicula consobrina</i> (Cailliaud, 1827)	+	-	+	+	-
<i>Pisidium pirothi</i> (Jickeli, 1881)	+	-	+	+	+
<i>Eupera ferruginea</i> (Krauss, 1848)	-	-	+	-	-
Total	15	43	39	24	9

Acknowledgments

The current study was a part of the project entitled 'Environmental Monitoring and Fisheries Development of Lake Nasser- towards sustainable development', which was funded by the National Institute of Oceanography and Fisheries (NIOF) of Egypt. The physicochemical parameters measured in this study were provided by the chemistry laboratory, Freshwater and Lakes Division of NIOF.

REFERENCES

- Abd El-Monsef, H.; Smith, S.E. and Darwish, K. (2015).** Impacts of the Aswan High Dam after 50 years. *Water Resour. Manag.*, **29**(6):1873–1885.
- Abdel-Gawad, S.S. (2016).** Impact of some physico-chemical factors of water on the diversity and distribution of Mollusca in Lake Nasser (Egypt). *Egypt. J. Aquat. Biol. & Fish.*, **20**(3): 89- 101.
- Abdel-Gawad, S.S. and Abdel-Aal, E.I. (2018).** Impact of Flood Cycle on Phytoplankton and Macroinvertebrates Associated with *Myriophyllum spicatum* in Lake Nasser Khors (Egypt). *J. Biol. Sci.*, **18**: 51- 67.
- Abdel-Gawad, S.S. and Mola, H.R. (2014).** Macrobenthic invertebrates in the main channel of Lake Nasser, Egypt. *Egypt. J. Aquat. Res.*, **40**: 405- 414.
- Abdo, M.H. (2003).** Physico-chemical characteristic of Abu Zabaal ponds, Egypt. *Egypt. J. Aquat. Res.*, **31** (2): 1- 15.
- Abou El-Ella, S.M. and El Samman, A. (2010).** Ecosystem status of the north part of Lake Nubia African. *J Biol Sci.*, **6**(2):7–21.
- Amer, A. S. (2007).** Effect of Different Types of Pollutants on Bacteria-Zooplankton Interaction in the Nile Water. Ph.D. Thesis, Fac. of Science, Zoology dep., Girls College, Ain Shams University. 227pp.
- APHA (2005).** American Water Works Association, Water Environment Federation. Standard Methods for the Examination of Water and Wastewater, 21th ed. American Public Health Association, Washington, DC. 2005.
- Brinkhurst, R.O. (1974).** The Benthos of Lakes. St. Martins Press, New York
- Elewa, A.A. (1987).** Fish stock assessment in Lake Nasser. A Nile Project, Inst. Oceanogr. and Fish., Academy of Scientific Research and Technology, Cairo, A.R.E.
- El-Shabrawy, G.M. (2001).** Ecological studies on Rotifera in Lake Qarun. *J. Egypt. Acad. Soc. Environ. Develop.* Vol, **2**(2): 1-18.
- El-Shabrawy, G.M. and Abd El-Regal, R. (1999).** Benthic fauna and sediment of Lake Nasser. I – Main channel and its littoral area. *Bulletin of the Faculty of Sciences, Zagazig University*, **21**: 193–215.

- El Gamal, T. and Zaki, N. (2017).** Egyptian irrigation after the Aswan High Dam. *Irrigated Agriculture in Egypt*, 47–79
- El Shemy, M. (2010).** Water quality modeling of large reservoirs in semiarid regions under climate change—example Lake Nasser, Egypt. Dissertation, University of Echnische.
- Entz, B. A. G. (1978).** Sedimentation processes in the Reservoir Lake Nasser -Nubia during 1965 -1974 and future aspects. Symposium of Ecological studies on the Nile River. Egyptian Society of Environment, Cairo.
- Farhat, H.I. and Aly, W. (2018).** Effect of site on sedimentological characteristics and metal pollution in two semi-enclosed embayments of great freshwater reservoir: Lake Nasser, Egypt. *J Afr Earth Sci.*, **141**: 194–206.
- Fishar, M.R.A. (1995).** Studies on bottom fauna in Lake Nasser, Egypt. Ph.D. Thesis, Suez Canal Univ., 267pp. 785
- Fishar, M.R.A. (2000).** Macrobenthic fauna, pp. 75–79. In J.F. Craig (ed.), *Sustainable Fish Production in Lake Nasser: Ecological Basis and Management Policy*. ICLARM Conference Proceedings, **61**: 184 pp.
- Gannon, J. E., & Stemberger, R. S. (1978).** Zooplankton (especially crustaceans and rotifers) as indicators of water quality. *Transactions of the American Microscopical Society*, 16-35.
- Hegab, M. H.; Khalifa, N., & Aly, W. (2021).** Zooplankton communities in Lake Nasser, Egypt, under the current flood regime, before the construction of Grand Ethiopian Renaissance Dam (GERD). *African Journal of Aquatic Science*, **46**(3), 329-339.
- Iskaros, I.A. (1988).** Biological studies on bottom fauna in Lake Nasser and adjacent waters. M.Sc. Thesis, Fac. Sci, Alexandria Univ., 184 pp.
- Iskaros, I.A. (1993).** Ecological studies on the distribution of zooplankton and bottom fauna in KhorKalabsha in Lake Nasser. Ph.D. Thesis, Fac. Sci, Mansoura Uni., 304 pp.
- Iskaros, I.A.; Gindy, N.N. and El Dardir, M. (2011).** Long-term fluctuations of macrobenthic invertebrates in Aswan Water Reservoir, Egypt. *Inter. J. Environ. Sci. Eng.* Vol.1: 37-48
- Iskaros, I.A. and El Dardir, M. (2010).** Factors affecting the distribution and abundance of bottom fauna in Lake Nasser, Egypt. *Nat. Sci.*, **8** (7): 95–110.
- Iskaros, I.A. and Gindy, N.N. (2009).** The effect of the substrate status on bottom fauna in Aswan reservoir, Egypt. *Egypt. J. Aquat. Res.*, **35**(2): 137-148.
- Khalifa, N.; El-Damhogy, K.A.; Fishar, M.R.; Nasef, A.M. and Hegab, M.H. (2015).** Using zooplankton in some environmental biotic indices to assess water quality of Lake Nasser, Egypt. *Int J Fisheries Aquat Stud.*, **2**(4): 281–289
- Latif, A.F.A. (1974).** Fisheries of Lake Nasser and Lake Nubia. In: Report on Trip to Lake Nasser and Lake Nubia, by Entz, B. and Latif, A.F.A. Aswan regional planning. Lake Nasser development center, A.R.E., pp.64-137.
- Latif, A.F.A.; El-Etreby, S.G.; Azim, M.E. and Sherie, H.A. (1979).** Reservoir fishes and fisheries resources. Report on Surveys to Lake Nasser and River Nile Project. Academy of Scientific Research and Technology, Cairo, Egypt.
- Mageed, A.A. and Heikal, M.T. (2006).** Factors affecting seasonal patterns in epilimnion zooplankton community in one of the largest man-made lakes in Africa (Lake Nasser, Egypt). *Limnologica*, **36**: 91- 97.
- Mola, H.R. and Abdel-Gawad, S.S. (2014).** Spatio-temporal variations of macrobenthic fauna in Lake Nasser khors, Egypt. *Egypt. J. Aquat. Res.*, **40**: 415- 423.

- Nassif, M.G. (2012).** Ecological studies on aquatic invertebrates of Ismailia Canal, Egypt. Ph.D. thesis, Fac. Science, Ain Shams Uni., 199 pp.
- Nkwoji, J. A.; Yakub, A.; Ajani, G. E.; Balogun, K. J.; Renner, K. O.; Igbo, J. K.; Ariyo, A. A. and Bello, B. O. (2010).** Seasonal variations in the water chemistry and benthic macroinvertebrates of a south western lagoon, Lagos, Nigeria. *Journal of American Science*. Vol. 6 (3): 85-92.
- Ravera, O. (1998).** Utility and limits of biological and chemical monitoring of the aquatic environment. *Annal.Dichim.*, **88**: 909-913. 786.
- Ravera, O. (2000).** Ecological monitoring for water body management. Proceedings of monitoring Tailormade III. International Workshop on Information for Sustainable Water Management, pp:157-167.
- Rosenberg, D.M.; Bilyj, B. and Wiens, A.P. (1984).** Chironomidae (Diptera) emerging from the littoral zone of reservoirs, with special reference to southern Indian Lake, Manitoba. *Can. J. Fish. Aquat. Sci.*, **41**: 638-647.
- Salem, T. A. (2021).** Changes in the physicochemical and biological characteristics in the lentic and lotic waters of the Nile river. *The Egyptian Journal of Aquatic Research*, *47*(1), 21-27.
- Salih, S.Q.; Allawi, M.F.; Yousif, A.A.; Armanuos, A.M.; Saggi, M.K.; Ali, M. and Chau, K.W. (2019).** Viability of the advanced adaptive neuro-fuzzy inference system model on reservoir evaporation process simulation: case study of Nasser Lake in Egypt. *EngApplComput Fluid Mech.*, **13**(1):878–891
- Wahab, H.M.A.; Iskaros, I.A.; Fishar, M.R. and Mohamed, N.E. (2018).** Ecological studies on macrobenthic invertebrates in four northern Khors of Lake Nasser (Egypt); Community Structure, Relative Abundance and Diversity. *Egypt. J. Aquat. Biol. & Fish.*, **22**(3): 111- 124.
- World Health Organization (WHO) (2003).** pH in drinking-water. Background document for preparation of WHO Guidelines for drinking-water quality. Geneva, World Health Organization (WHO/SDE/WSH/03.04/12).
- Yousry, K.M. (2009).** Assessment and Modeling of Water Quality and Zooplankton of El-Salam Canal, Egypt. Ph. D. Thesis, Dep. Environmental Basic Sciences. Institute of Environmental Studies and Research. Ain Shams University.