



## The Second Study on Zooplankton in the Tigris River in Misan Province, Iraq Since 1921: Evidence to Include Zooplankton in Ecosystem Monitoring

Ahmed S. Mohammed Al-Budeiri<sup>1</sup>, Ghassan A. Ali Al-Yacoub<sup>2\*</sup>, Firas S. Idan<sup>3</sup>

<sup>1</sup>Department of Biology, College of Science, University of Misan, Misan, Iraq

<sup>2</sup>Department of Biology, College of Education for Pure Sciences, University of Thi-Qar, Thi-Qar, 64001, Iraq

<sup>3</sup>General Directorate of Education for Misan Governorate, Ministry of Education, Iraq

\*Corresponding author: ghassanadnan.bio@utq.edu.iq

### ARTICLE INFO

#### Article History:

Received: July 16, 2024

Accepted: Aug. 5, 2024

Online: Sep. 1<sup>st</sup>, 2024

#### Keywords:

Cladocera,  
Copepoda,  
Environmental  
monitoring,  
Iraq,  
Rotifera,  
Tigris River,  
Zooplankton

### ABSTRACT

The present study was conducted in one of the most significant water sources in Iraq, the Tigris River in Misan Province. The aim was to examine the zooplankton community, specifically Cladocera, Copepoda, and Rotifera. Samples were collected 30cm below the water surface using a plankton haul net with a mesh size of 64µm from September 2023 to February 2024. Sampling took place at three stations: the first in Kumait, located in the north of Misan Province; the second in Amarah City, the capital of Misan Province, and the third about 20km south of Amarah City. The results revealed seventeen species of zooplankton, belonging to two phyla, three classes, four orders, and eight families. A comparison between the historical and current status in the study area indicated a noticeable shift in zooplankton composition toward species more tolerant of pollution. For instance, the appearance of the family Brachionidae and its common species *Brachionus* sp. in the study area, alongside the absence of *Daphnia* species, suggests ecological degradation in the inland waters of Iraq, particularly in Misan Province. This study recommends incorporating zooplankton into water quality monitoring programs (WQMPS) in Iraq.

### INTRODUCTION

Zooplankton are important groups in aquatic ecosystems (i.e. rivers, lakes, swamps and oceans) due to their central position in the food webs of these systems (Persson & Vrede, 2006). The productions of the lower food webs components which are mainly derived from algae, bacteria and protozoa are mostly linked with higher trophic levels (e.g. macroinvertebrates and fish) via zooplankton (Semyalo *et al.*, 2009). Thus, they play a crucial role in the energy flow through aquatic food webs and in maintaining the

efficiency of these organisms (Azam *et al.*, 1983; Vadstein *et al.*, 1989; Arndt, 1993). In addition, zooplankton play an important role in nutrient recycling through the remineralization of essential nutrients such as N and P, which can be taken by algae and bacteria, and thus encouraging their growth (Walve & Larsson, 1999; Likens, 2010). Due to their small size, short lifespans, and sensitivity to different pollutants and impacts of climate change, zooplankton composition serves as an indicator for water quality, helping to determine the ecological state of different aquatic ecosystems (Wetzel, 2001; Lampert & Sommer, 2007; Jeppesen *et al.*, 2011; Chiba *et al.*, 2018; Labuce *et al.*, 2020).

The most important components of zooplankton are Rotifera, Copepoda and Cladocera. Rotifera are an essential component of the aquatic food webs (Gilbert, 2022). Rotifers are composed of 2000 species (Ricci & Balsamo, 2000). Most rotifers can be found in freshwater, but about 100 species live in brackish or marine habitats (Likens, 2010). They comprise about 30% of the biomass of freshwater plankton in some habitats (Nogrady *et al.*, 1993). They are more important consumers in feeding on small phytoplankton and components of the microbial loop (i.e. bacteria, ciliates and flagellates) in freshwater ecosystems in comparison with copepods and cladocerans (Thorp & Covich, 2009).

Cladocera are crustaceans found in fresh, brackish and marine water, found in permanent and temporary water bodies. Cladocerans are considered as filter-feeders (Forro *et al.*, 2008), feeding on bacteria, phytoplankton and protozoans. The numbers of species of Cladocera are less than 600, and only 2% live in marine ecosystems (Dumont & Negrea, 1996). Typically, their size ranges from 0.2 to 6mm (Forró *et al.*, 2008). Additionally, it is also reported that their size can reach 18mm in species such as *Leptodora kindtii* (Fryer, 1987).

Crustacean copepods live in fresh and hypersaline water habitats with about 900 species and form a significant part of plankton communities (Dole-Olivier *et al.*, 2000; Mondal *et al.*, 2013). The copepods are selective feeders (Djeghri *et al.* 2018). They mainly feed on bacteria (Wroblewski, 1980; Zöllner *et al.*, 2003). Besides, they feed on phytoplankton (Calbet *et al.*, 2000). Thus, they serve as an important trophic link between the bases of the food web with higher trophic levels (Verity & Smetacek, 1996).


The first study on the composition of zooplankton in Iraq was conducted by Gurney (1921), who studied zooplankton around Amara City near the Tigris River in Misan Province during the period between 1917 and 1918. Gurney (1921) recorded 9 species of Cladocera, 4 species of Copepoda and 3 species of Rotifera. These species which were mainly collected from permanent marsh and temporary pools close to the Tigris River were approximately representative of the zooplankton community in the river. To the author knowledge, no studies have been conducted on zooplankton in this part of the

Tigris River since 1921. The aim of this study was to provide the second information about zooplankton composition in this area.

## MATERIALS AND METHODS

### 1. Study area

The Tigris River was an important river for people during Mesopotamian civilization (Varol *et al.*, 2010), and is considered to be one of the main water sources in addition to the Euphrates River in Iraq (Al-Ansari *et al.*, 2019). The Tigris River primarily originates from Turkey. The length of the river is 1900km, and most of the river length is located in Iraq with a percentage of approximately 77% by about 1415km of its actual length (Kibaroglu, 2002). The part of the river in Misan Province is about 160km long. The map was prepared using SimpleMappr (Shorthouse, 2010) (Fig. 1). The first station is located to the north of Misan Province, called Kumait ( $32^{\circ}02'00.6''\text{N}$   $46^{\circ}52'49.5''\text{E}$ ); the second station is approximately in the center of Amarah City ( $31^{\circ}45'21.2''\text{N}$   $47^{\circ}09'00.5''\text{E}$ ), and the third station is located to the south of Amarah City by about 20km from the last station ( $31^{\circ}41'16.7''\text{N}$   $47^{\circ}09'56.8''\text{E}$ ). Several stations were chosen to collect representative samples. The second and three stations receive huge amounts of untreated wastewater discharges from households and hospitals in Amarah City. The population of Misan Province is about 1.025.862 (NIC 2023), and most of them living in Amarah City.



**Fig. 1.** Sample collection stations on the Tigris River in Misan Province, Kumait district (Black circle), Amarah City (Red circle), south of Amarah City (Yellow circle)

## 2. Collection of zooplankton samples

Zooplankton samples were collected from September 2023 to February 2024 from three stations on the Tigris River in Misan Province.

Samples were collected 30cm below the water surface using a plankton haul net with a mesh size of 64µm. After each sampling, the net was washed to avoid potential contamination of gathered samples from previous sampling (Goswami, 2004). Then, they were preserved in 70% ethanol, and all preserved samples were transferred back to the laboratory. A Sedgewick Rafter chamber and Olympus compound Microscope were used for classifying zooplankton. Identification of zooplankton groups (Cladocera, Copepoda and Rotifera) was based on the taxonomic keys of Edmondson (1959), Korinek (1999) and Fernando (2003). Zooplankton were classified to species level.

## RESULTS AND DISCUSSION

In this study seventeen species of zooplankton were recorded in the Tigris River in Misan Province which are shown in Tables (1, 2, 3). These species belong to two phyla, three classes, four orders and eight families. Station one was more diverse in its species composition than other stations. This may be due to these stations receiving direct high discharge of wastewater without treatment. Many studies on the Tigris River and in its part in Misan Province (Amteghy, 2014; Al-Mussawy *et al.*, 2023; Al-Budeiri *et al.*, 2024; Hameed, 2024) have reported high levels of pollutants in this river. These studies indicated that lacking wastewater treatment was the main reason. It is now known that the zooplankton community structure is affected by pollution, thus zooplankton species are considered to be indicators for assessing the environmental condition (Singh & Sharma, 2020; Boldrocchi *et al.*, 2023). Jeppesen *et al.* (2011) showed that zooplankton are significant biological quality element and should be included as a central element in aquatic monitoring programs of the EU.

Gurney (1921) who studied zooplankton in Amara City between 1917 and 1918, recorded nine species of cladocerans, and four species of Copepoda, and three taxa of rotifers (Table 4). The comparison between data on the species composition of Gurney (1921), and data about zooplankton species presented here in this study (Table 4) showed a change in the species of zooplankton community, suggesting that the current ecological status in the study area is unfavorable for most species of zooplankton which is recorded by Gurney (1921). Recently, the major decline in environmental condition of the Tigris River due to the impacts of pollution and climate change associated with high temperatures and low precipitation (Al-Saady & Abdullah, 2014; Abbas *et al.*, 2016; Price, 2018; Al-Saedi *et al.*, 2024). This is expected to be the main drivers for such sharp change in species composition of zooplankton (Sterza & Fernandes, 2006; Jeppesen *et al.*, 2015), and in the study area. For example, many species recorded by Gurney (1921), such as *Daphnia lumholtzi*, *Daphnia longispina*, *Scapholeberis* sp., *Ceriodaphnia* sp.,

and *Alona* sp. (Greenwald & Hurlbert, 1993; Gonçalves *et al.*, 2007; Dao *et al.*, 2018), are specialized for freshwater habitats and are less tolerant to chemical and biological pollutants, such as pesticides, increased salinity, heavy metals, and toxins produced by cyanobacteria. These pollutants have become common in Iraqi waters and worldwide (DouAbui *et al.*, 1988; Le, 2020; Gummaa *et al.*, 2021; Hmoshi & Mohammed, 2022; Hameed, 2024). As a result, these zooplankton, particularly *Daphnia* species, are frequently used in aquatic biomonitoring due to their high sensitivity to pollution (Tomasiks & Warren, 1996; Le *et al.*, 2016; Abdullahi *et al.*, 2022). Zooplankton can serve as bioindicators based on species-level changes (De Eyto *et al.*, 2003). The zooplankton species recorded in this study, such as *Anuraeopsis fissa*, *Bosmina longirostris*, *Bosmina meridionalis*, *Brachionus angularis*, *Brachionus plicatilis*, *Chydorus sphaericus*, *Eucyclops agilis*, *Filinia longiseta*, *Mesocyclops edax*, *Notholca acuminata*, *Simocephalus vetulus*, and *Trichocerca longiseta*, along with the first occurrence of the family Brachionidae and its common species *Brachionus* sp., indicate low water quality in the study area (De Eyto *et al.*, 2003; Ezz *et al.*, 2014; Costa *et al.*, 2016; Perbiche-Neves *et al.*, 2016; Al-Kalidy *et al.*, 2017; Loria, 2017; Leppänen, 2018; Muñoz-Colmenares *et al.*, 2021; de Paiva *et al.*, 2023).

The tolerance or sensitivity of zooplankton species for disturbances in aquatic ecosystem or for polluted water are mainly dependent on their range of responses, adaptations and evolution to ecological changes (Thackeray & Beisner, 2024). These changes, for example, could appear in seasonal cycle, thus, zooplankton will show patterns of seasonality in their structure (Salman *et al.*, 2014; Ajeel & Abbas, 2019; Van Engeland *et al.*, 2023), if ecological changes last for decades or for centuries with ongoing dramatic increases in the levels and effects of these changes due to anthropogenic activities combined with climate change impacts; this will cause a shift or changes in zooplankton community (Marques *et al.*, 2024; Pershing & Kemberling, 2024).

The changes in zooplankton composition will ultimately have huge impact on structure and function of aquatic ecosystems (Nielsen *et al.*, 2003).

**Table 1.** Zooplankton composition in the Tigris River in Misan Province at station 1

Zooplankton Groups	Taxon
Phylum Arthropoda	
Subphylum Crustacea	
Class Branchiopoda	
Order Cladocera	
1-Family Bosminidae	<i>Bosmina longirostris</i> Müller, 1776
2- Family Chydoridae	<i>Chydorus sphaericus</i> Müller, 1776
3- Family Daphniidae	<i>Simocephalus vetulus</i> Schoedler, 1858
Class Maxillopoda	
Subclass Copepoda	
Order Cyclopoida	
1-Family Cyclopidae	<i>Mesocyclops edax</i> Forbes, 1891
	<i>Cyclops scutifer</i> Sars, 1863
	<i>Eucyclops agilis</i> Koch, 1838
	<i>Orthocyclops modestus</i> Herrick, 1883
Phylum Rotifera	
Class Monogononta	
Order Flosculariaceae	
1-Family Trochosphaeridae	<i>Filinia longiseta</i> Ehrenberg, 1834
Order Ploima	
2-Family Brachionidae	<i>Anuraeopsis fissa</i> Gosse, 1851
	<i>Brachionus angularis</i> Gosse, 1851
	<i>Brachionus rotundiformis</i> Tschugunoff, 1921
	<i>Notholca acuminata</i> Ehrenberg, 1832
3- Family Euchlanidae	<i>Euchlanis deflexa</i> Gosse, 1851

4-Family Trichocercidae	<i>Trichocerca longiseta</i> Schrank, 1802
-------------------------	--------------------------------------------

**Table 2.** Zooplankton composition in the Tigris River in Misan Province at station 2

Zooplankton Groups	Taxon
Phylum Arthropoda	
Subphylum Crustacea	
Class Branchiopoda	
Order Cladocera	
1-Family Bosminidae	<i>Bosmina longirostris</i> Müller, 1785
2- Family Chydoridae	<i>Chydorus sphaericus</i> Müller, 1776
Class Maxillopoda	
Subclass Copepoda	
Order Cyclopoida	
1-Family Cyclopidae	<i>Mesocyclops edax</i> Forbes, 1891
	<i>Cyclops scutifer</i> Sars, 1863
	<i>Diacyclops thomasi</i> Forbes, 1882
Phylum Rotifera	
Class Monogononta	
Order Ploima	
1-Family Brachionidae	<i>Brachionus angularis</i> Gosse, 1851
	<i>Brachionus rotundiformis</i> Tschugunoff, 1921

**Table 3.** Zooplankton composition in the Tigris River in Misan Province at station 3

Zooplankton Groups	Taxon
Phylum Arthropoda	
Subphylum Crustacea	
Class Branchiopoda	
Order Cladocera	
1-Family Bosminidae	<i>Bosmina longirostris</i> Müller, 1776
	<i>Bosmina Meridionalis</i> Sars, 1904

2- Family Chydoridae	<i>Chydorus sphaericus</i> Müller, 1776
Class Maxillopoda	
Subclass Copepoda	
Order Cyclopoida	
1-Family Cyclopidae	<i>Mesocyclops edax</i> Forbes, 1891
	<i>Cyclops scutifer</i> Sars, 1863
Phylum Rotifera	
Class Monogononta	
Order Ploima	
1-Family Brachionidae	<i>Brachionus angularis</i> Gosse, 1851
	<i>Brachionus rotundiformis</i> Tschugunoff, 1921
	<i>Brachionus plicatilis</i> Müller, 1786

**Table 4.** Zooplankton species recorded in the present study and by **Gurney (1921)**

Zooplankton Groups	Taxon recorded in the present study	Taxon recorded by Gurney (1921)
Phylum Arthropoda		
Subphylum Crustacea		
Class Branchiopoda		
Order Cladocera		
1-Family Bosminidae	<i>Bosmina longirostris</i> Müller, 1776	Present
	<i>Bosmina Meridionalis</i> Sars, 1904	Not present
2- Family Chydoridae	<i>Chydorus sphaericus</i> Müller, 1776	Present
	<i>Alona rectangular</i> Sars, 1862	Not present
	<i>Alona costata</i> Sars, 1862	Not present
3- Family Daphniidae	<i>Simocephalus vetulus</i> Schoedler, 1858	Not present
	Not present	<i>Ceriodaphnia reticulata</i>



		Jurine, 1820
	Not present	<i>Daphnia lumholtzi</i> Sars, 1885
	Not present	<i>Daphnia longispina</i> Müller, 1776
	Not present	<i>Scapholeberis mucronate</i> Müller, 1776
	Not present	<i>Simocephalus exspinosus</i> De Geer, 1778
Class Maxillopoda		
Subclass Copepoda		
Order Cyclopoida		
1-Family Cyclopidae	<i>Mesocyclops edax</i> Forbes, 1891	Not present
	<i>Cyclops scutifer</i> Sars, 1863	Not present
	<i>Eucyclops agilis</i> Koch, 1838	Present
	<i>Orthocyclops modestus</i> Herrick, 1883	Not present
	<i>Diacyclops thomasi</i> Forbes, 1882	Not present
	Not present	<i>Cyclops vicinus</i> Uljanin, 1875
Order Calanoida		
1-Family Diaptomidae	Not present	<i>Diaptomus vulgaris</i> Schmeil, 1896
Order Harpacticoida		
1-Family Canthocamptidae	Not present	<i>Canthocamptus staphylinus</i> Jurine, 1820
Phylum Rotifera		
Class Monogononta		
Order Flosculariaceae		

1-Family Trochosphaeridae	<i>Filinia longiseta</i> Ehrenberg, 1834	Not present
Order Ploima		
2-Family Brachionidae	<i>Anuraeopsis fissa</i> Gosse, 1851	Not present
	<i>Brachionus angularis</i> Gosse, 1851	Not present
	<i>Brachionus rotundiformis</i> Tschugunoff, 1921	Not present
	<i>Brachionus plicatilis</i> Müller, 1786	Not present
	<i>Notholca acuminata</i> Ehrenberg, 1832	Not present
3- Family Euchlanidae	<i>Euchlanis deflexa</i> Gosse, 1851	Not present
4-Family Trichocercidae	<i>Trichocerca longiseta</i> Schrank, 1802	Not present
5- Family Trichotriidae	Not present	<i>Dinocharis pocillum</i> , Müller, 1776
6- Family Gastropodidae	Not present	<i>Ascomorpha</i> sp. Perty, 1850
7- Family Asplanchnidae	Not present	<i>Asplanchna</i> sp. Gosse, 1850

## CONCLUSION

The comparison between the historical and present status of the study area reveals a noticeable shift in zooplankton composition toward species that are more tolerant of pollution. For example, the emergence of the family Brachionidae and its common species, *Brachionus* sp., and the absence of *Daphnia* species in the current study indicate ecological degradation in the inland waters of Iraq. This study suggests that zooplankton should be included in water quality monitoring programs (WQMPs) in Iraq. More

comprehensive studies are needed, employing a wide range of ecological tools, to investigate the responses of zooplankton to different stressors.

## REFERENCES

- Abbas, N.; Wasimi, S.A. and Al-Ansari, N.** (2016). Climate change impacts on water resources of Greater Zab River, Iraq. *Journal of Civil Engineering and Architecture*, 10(12), 1384-1402.
- Al-Ansari, N.; Jawad, S.; Adamo, N. and Sissakian, V.** (2019). Water quality and its environmental implications within Tigris and Euphrates rivers. *Journal of Earth Sciences and Geotechnical Engineering*, 9(4), 57-108.
- Al-Budeiri, A.S.M.; Idan, F.S.; Al-Yacoub, G.A. Ali and Jazza, S.H.** (2024). Some Ecological Characteristics of Um-Al-Naaj Marsh, Southern Iraq in the Second Most Driest Season during the Last 40 Years. *EJABF*, 28(3): 1343 – 1354. <https://doi.org/10.21608/ejabf.2024.362732>.
- Al-Kalidy, S. K. A. A.; AL-Keriawy, H. A. H. and Ahmod, Q. A.** (2017). Qualitative and Quantitative composition of Rotifers community in Eastern Euphrates drainage canal/Al-Qadisiyha province. *J. Euphrates Journal of Agriculture Science*, 9(2), 13-27.
- Al-Mussawy, H.A.; Mohamed, N.H. and Al-Madhhachi, A.S.T.** (2023). Monitoring the water quality of the Tigris River for drinking and irrigation purposes in Maysan Province, Iraq. *Sustain. Water Resour. Manag.*, 9(6), 177.
- Al-Saady, Y.I. and Abdullah, E.J.** (2014). Water Quality of Tigris River within Missan Governorate eastern part of the Mesopotamia Plain–Iraq. *JUB*, 22(9), 2489-2502.
- Al-Saedi, M.S.; Naimi, S. and Al-Sharify, Z.T.** (2024). The environmental impact of pollutants and heavy materials on the water quality in the Tigris River. *MMEP*, 11(2), 290-300. <https://doi.org/10.18280/mmep.110202>.
- Abdullahi, M.; Li, X.; Abdallah, M.A.E.; Stubbings, W.; Yan, N.; Barnard, M.; ... and Orsini, L.** (2022). *Daphnia* as a sentinel species for environmental health protection: a perspective on biomonitoring and bioremediation of chemical pollution. *Environ. Sci. Technol.*, 56(20), 14237-14248. <https://doi.org/10.1021/acs.est.2c01799>.
- Ajeel, S. G. and Abbas, M. F.** (2019). Diversity, Abundance, and Distribution of Cladocera at the end of the Tigris River North of Basrah–IRAQ. *Baghdad Science Journal*, 16(4).
- Amteghy, A.H.** (2014). Impact of Sewage Water on the Water Quality of Tigris River in Maysan Province and their Possible Health Risks and Removal by Using Granular Activated Carbon and Sand. *MJB*, 11(1).
- Azam, F.; Fenchel, T.; Field, J.G.; Gray, J.S.; Meyer-Reil, L.A. and Thingstad, F.** (1983). The ecological role of water-column microbes in the sea. *MEPS*, 10(3), 257-263. <https://doi.org/10.7208/chicago/9780226125534-024>.

- Boldrocchi, G.; Villa, B.; Monticelli, D.; Spanu, D.; Magni, G.; Pachner, J.; ... and Bettinetti, R.** (2023). Zooplankton as an indicator of the status of contamination of the Mediterranean Sea and temporal trends. *Mar. Pollut. Bull.*, 197, 115732. <https://doi.org/10.1016/j.marpolbul.2023.115732>.
- Calbet, A.; Landry, M.R. and Scheinberg, R.D.** (2000). Copepod grazing in a subtropical bay: Species-specific responses to a midsummer increase in nanoplankton standing stock. *MEPS*, 193, 75-84.
- Costa, B. N. S., Pinheiro, S. C. C., Amado, L. L. and de Oliveira Lima, M.** (2016). Microzooplankton as a bioindicator of environmental degradation in the Amazon. *Ecological Indicators*, 61, 526-545.
- Dao, T.S.; Wiegand, C.; Bui, B.T. and Dinh, K.V.** (2018). Transgenerational effects of cyanobacterial toxins on a tropical micro-crustacean *Daphnia lumholtzi* across three generations. *Environ. Pollut.*, 243, 791-799. <https://doi.org/10.1016/j.envpol.2018.09.055>.
- De Eyto, E.; Irvine, K.; García-Criado, F.; Gyllström, M.; Jeppensen, E.; Kornijow, R.; ... and Moss, B.** (2003). The distribution of chydorids (Branchiopoda, Anomopoda) in European shallow lakes and its application to ecological quality monitoring. *Archiv für Hydrobiologie*, 156(2), 181-202.
- de Paiva, L. P.; Cavalcanti, V. B.; Soares Filho, A. A.; Viana, W. K. R.; da Silva Apoliano, M. L. and Souza, R. L. M.** (2023). Water quality and phyto-zooplanktonic bioindicators in the cocó river estuary, fortaleza/ce. *Revista Contemporânea*, 3(11), 20833-20848.
- Djehri, N.; Atkinson, A.; Fileman, E.S.; Harmer, R.A.; Widdicombe, C.E.; McEvoy, A.J.; ... and Mayor, D.J.** (2018). High prey-predator size ratios and unselective feeding in copepods: a seasonal comparison of five species with contrasting feeding modes. *Prog. Oceanogr.*, 165, 63-74. <https://doi.org/10.1016/j.pocean.2018.04.013>.
- Dole-Olivier, M.J.; Galassi, D.M.P.; Marmonier, P. and Creuzé des Châtelliers, M.** (2000). The biology and ecology of lotic microcrustaceans. *Freshw. Biol.*, 44(1), 63-91. <https://doi.org/10.1046/j.1365-2427.2000.00590.x>.
- DouAbui, A.A.; Al-Saad, H.T.; Al-Timari, A.A. and Al-Rekabi, H.N.** (1988). Tigris-Euphrates Delta: a major source of pesticides to the Shatt al-Arab River (Iraq). *AECT*, 17, 405-418. <https://doi.org/10.1007/BF01055178>.
- Dumont, H.J. and Negrea, S.** (1996). A conspectus of the cladocera of the subterranean waters of the world. *Hydrobiologia*, 325(1), 1-30.
- Edmondson, W.T.** (1959). *Freshwater Biology*. John Wiley and Sons Inc., New York, 1248 p.
- Ezz, S. M. A.; Aziz, N. E. A.; Abou Zaid, M. M.; El Raey, M. and Abo-Taleb, H. A.** (2014). Environmental assessment of El-Mex Bay, Southeastern Mediterranean by

- using Rotifera as a plankton bio-indicator. The Egyptian Journal of Aquatic Research, 40(1), 43-57.
- Fernando, C.H.** (2003). A guide to tropical freshwater zooplankton: identification, ecology and impact on fisheries. QRB, 78(2), 291pp.
- Forró, L.; Korovchinsky, N.M.; Kotov, A.A. and Petrusek, A.** (2008). Global diversity of cladocerans (Cladocera; Crustacea) in freshwater. Hydrobiologia, 595, 177–184. <https://doi.org/10.1007/s10750-007-9013-5>.
- Fryer, G.** (1987). Morphology and the classification of the so-called Cladocera. Hydrobiologia, 145(1), 19-28.
- Gilbert, J.J.** (2022). Food niches of planktonic rotifers: Diversification and implications. L&O, 67(10), 2218-2251. <https://doi.org/10.1002/lno.12199>.
- Gonçalves, A.M.M.; Castro, B.B.; Pardal, M.A. and Gonçalves, F.** (2007). Salinity effects on survival and life history of two freshwater cladocerans (*Daphnia magna* and *Daphnia longispina*). In Annales De Limnologie- IJLI, 43, (1), 13-20. <https://doi.org/10.1051/limn/2007022>.
- Goswami, S.C.** (2004). Zooplankton methodology, collection and identification - A field manual. National Institute of Oceanography (NIO), 26.
- Greenwald, G.M. and Hurlbert, S.H.** (1993). Microcosm analysis of salinity effects on coastal lagoon plankton assemblages. Hydrobiologia, 267, 307-335.
- Gummaa, N.R.; Dwaish, A.S. and Hamzah, I.H.** (2021). Molecular detection of some toxogenic cyanobacteria in Tigris River in Baghdad–Iraq. Mol. Biol. Rep., 48(7), 5393-5397. <https://doi.org/10.1007/s11033-021-06538-z>.
- Gurney, R.** (1921). Freshwater Crustacea collected by Dr. P. A. Buxton in Mesopotamia and Persia. JBNHS, 27, 835-843.
- Hameed, A.A.S.** (2024). Pollution of Water's direct effect in Iraq on the Public Health & Safety. JFES, 2(2), 12-12.
- Hmoshi, R.M. and Mohammed, M.I.** (2022). Cyanobacteria diversity in various waterbodies of Mosul, Iraq. Int. J. Aquat. Biol., 10(6), 537-542.
- Jeppesen, E.; Brucet, S.; Naselli-Flores, L.; Papastergiadou, E.; Stefanidis, K.; Nöges, T.; ... and Beklioğlu, M.** (2015). Ecological impacts of global warming and water abstraction on lakes and reservoirs due to changes in water level and related changes in salinity. Hydrobiologia, 750, 201-227. <https://doi.org/10.1007/s10750-014-2169-x>.
- Jeppesen, E.; Nöges, P.; Davidson, T.A.; Haberman, J.; Nöges, T.; Blank, K.; ... and Amsinck, S.L.** (2011). Zooplankton as indicators in lakes: a scientific-based plea for including zooplankton in the ecological quality assessment of lakes according to the European Water Framework Directive (WFD). Hydrobiologia, 676, 279-297. <https://doi.org/10.1007/s10750-011-0831-0>.
- Kibaroglu, A.** (2002). Building a regime for the waters of the Euphrates-Tigris river basin. Kluwer Law International, 319pp.

- Korinek, V.** (1999). A guide to limnetic species of Cladocera of African inland Waters (Crustacea, Brachiopoda) (Using the morphology of parthenogenetic females). Volta Basin Research Project, Accra on behalf of International Association of Theoretical and Applied Limnology c/o Department of Biological Sciences, University of Alabama, Tuscaloosa, Alabama.
- Labuce, A.; Dimante-Deimantovica, I.; Tunens, J. and Strake, S.** (2020). Zooplankton indicator-based assessment in relation to site location and abiotic factors: A case study from the Gulf of Riga. *Environ. Monit. Assess.*, 192(2), 147. <https://doi.org/10.1007/s10661-020-8113-9>.
- Lampert, W. and Sommer, U.** (2007). *Limnology: the ecology of lakes and streams*. Oxford University Press, USA.
- Le, T.D.H.** (2020). Pesticides and salinisation, two stressors of freshwater ecosystems. PhD thesis, University of Koblenz-Landau, 149pp.
- Le, Q.A.V.; Sekhon, S.S.; Lee, L.; Ko, J.H. and Min, J.** (2016). *Daphnia* in water quality biomonitoring-“omic” approaches. *Toxicol. Environ. Health Sci.*, 8, 1-6. <https://doi.org/10.1007/s13530-016-0255-3>.
- Leppänen, J. J.** (2018). An overview of Cladoceran studies conducted in mine water impacted lakes. *International Aquatic Research*, 10(3), 207-221.
- Likens, G. (Ed.).** (2010). *Plankton of inland waters*. San Diego: CA: Academic Press/Elsevier.
- Loria, K.** (2017). Freshwater zooplankton communities as indicators of habitat quality: Testing responses to multiple disturbances. University of Colorado at Boulder. Undergraduate honors theses, 1388.
- Marques, R.; Otto, S. A.; Di Pane, J.; Boersma, M.; Meunier, C. L.; Wiltshire, K. H.; ... and Renz, J.** (2024). Response of the meso-and macro-zooplankton community to long-term environmental changes in the southern North Sea. *ICES Journal of Marine Science*, 81(3), 526-539.
- Mondal, D.; Pal, J.; Ghosh, T.K. and Biswas, A.K.** (2013). Diversity of cladocerans and copepods of Mirik Lake in Darjeeling himalaya. *JTBSSRR*, 2(1), 36-46.
- Muñoz-Colmenares, M. E.; Soria, J. M. and Vicente, E.** (2021). Can zooplankton species be used as indicators of trophic status and ecological potential of reservoirs?. *Aquatic Ecology*, 55(4), 1143-1156.
- National Investment Commission.** (2023). <https://investpromo.gov.iq/wp-content/uploads/2013/05/Maysan-province-En.pdf>. Accessed 15 February 2024.
- Nielsen, D.L.; Brock, M.A.; Rees, G.N. and Baldwin, D.S.** (2003). Effects of increasing salinity on freshwater ecosystems in Australia. *Aust. J. Bot.*, 51(6), 655-665. <https://doi.org/10.1071/BT02115>.
- Nogrady T.; Wallace R.L. and Snell T.W.** (1993). *Rotifera. Vol 1: Biology, Ecology and Systematics. Guides to the Identification of the Microinvertebrates of the Continental Waters of the World*. SPB Academic publishing, the Hague.

- Perbiche-Neves, G.; Saito, V. S.; Previattelli, D.; Da Rocha, C. E. and Nogueira, M. G.** (2016). Cyclopoid copepods as bioindicators of eutrophication in reservoirs: Do patterns hold for large spatial extents?. *Ecological Indicators*, 70, 340-347.
- Pershing, A. J. and Kemberling, A.** (2024). Decadal comparisons identify the drivers of persistent changes in the zooplankton community structure in the Northwest Atlantic. *ICES Journal of Marine Science*, 81(3), 564-574.
- Persson, J. and Vrede, T.** (2006). Polyunsaturated fatty acids in zooplankton: variation due to taxonomy and trophic position. *Freshw. Biol.*, 51(5), 887-900. <https://doi.org/10.1111/j.1365-2427.2006.01540.x>.
- Price, R.A.** (2018). Environmental risks in Iraq, K4D Helpdesk Report. Brighton, UK: Institute of Development Studies, 16pp.
- Ricci, C. and Balsamo, M.** (2000). The biology and ecology of lotic rotifers and gastrotrichs. *Freshw. Biol.*, 44(1), 15-28.
- Salman, S. D.; Abbas, M. F.; Ghazi, A. H. M.; Ahmed, H. K.; Akash, A. N.; Douabul, A. A.; ... and Asada, T.** (2014). Seasonal changes in zooplankton communities in the re-flooded Mesopotamian wetlands, Iraq. *Journal of freshwater Ecology*, 29(3), 397-412.
- Semyalo, R.; Nattabi, J.K. and Larsson, P.** (2009). Diel vertical migration of zooplankton in a eutrophic bay of Lake Victoria. *Hydrobiologia*, 635(1), 383-394. <https://doi.org/10.1007/s10750-009-9931-5>.
- Shorthouse, D.P.** (2010). SimpleMappr, an online tool to produce publication-quality point maps. Available from: <http://www.simplemappr.net> (accessed 16 March 2024).
- Singh, S. and Sharma, R.C.** (2020). Zooplankton diversity and potential indicator species for assessment water quality of high altitude wetland, Dodi Tal of Garhwal Himalaya. *India Academia Arena*, 12(5), 1-16. <https://doi.org/10.7537/marsaaj120520.01>.
- Sterza, J.M. and Fernandes, L.L.** (2006). Distribution and abundance of Cladocera (Branchiopoda) in the Paraíba do Sul river estuary, Rio de Janeiro, Brazil. *Braz. J. Oceanogr.*, 54, 193-204.
- Thackeray, S. J. and Beisner, B. E.** (2024). Zooplankton Communities: Diversity in Time and Space. In *Wetzel's Limnology* (pp. 539-585). Academic Press.
- Thorp, J.H. and Covich, A.P.** (2009). Ecology and classification of North American Freshwater Invertebrates, 2nd ed.; Academic press, 1005pp.
- Tomasiks, P. and Warren, D. M.** (1996). The use of *Daphnia* in studies of metal pollution of aquatic systems. *Environ. Rev.*, 4(1), 25-64.
- Vadstein, O.; Harkjerr, B.O.; Jensen, A.; Olsen, Y. and Reinertsen, H.** (1989). Cycling of organic carbon in the photic zone of a eutrophic lake with special reference to the heterotrophic bacteria. *L&O*, 34(5), 840-855.
- Van Engeland, T.; Bagøien, E.; Wold, A.; Cannaby, H. A.; Majaneva, S.; Vader, A.; ... and Ingvaldsen, R. B.** (2023). Diversity and seasonal development of large

zooplankton along physical gradients in the Arctic Barents Sea. Progress in Oceanography, 216, 103065.

**Varol, M.; Gökot, B. and Bekleyen, A.** (2010). Assessment of water pollution in the Tigris River in Diyarbakır, Turkey. Water pract. technol., 5(1), wpt2010021. <https://doi.org/10.2166/wpt.2010.021>.

**Verity, P.G. and Smetacek, V.** (1996). Organism life cycles, predation, and the structure of marine pelagic ecosystems. Mar. Ecol. Prog. Ser., 130: 277-293.

**Walve, J. and Larsson, U.** (1999). Carbon, nitrogen and phosphorus stoichiometry of crustacean zooplankton in the Baltic Sea: implications for nutrient recycling. J. Plankton Res., 21(12), 2309-2321.

**Wetzel R.G.** (2001). Limnology: Lake and River Ecosystems, 3rd ed. Academic Press, San Diego, 1006pp.

**Wroblewski, J.** (1980). A simulation of the distribution of *Acartia clausi* during Oregon upwelling, august 1973. J. Plankton Res., 2(1), 43-68.

**Zöllner, E.; Santer, B.; Boersma, M.; Hoppe, H. and Jürgens, K.** (2003). Cascading predation effects of *Daphnia* and copepods on microbial food web components. Freshw. Biol., 48(12), 2174-2193. <https://doi.org/10.1046/j.1365-2426.2003.01158.x>.